

Canaries in a Coal Mine: Using Globular Clusters to Place Limits on Massive Black Holes in the Galactic Halo

Phil Arras and Ira Wasserman

Center for Radiophysics and Space Research, Cornell University, Ithaca, NY 14853

1 February 2008

ABSTRACT

We explore the possibility that massive black holes comprise a significant fraction of the dark matter of our galaxy by studying the dissolution of galactic globular clusters bombarded by them. In our simulations, we evolve the clusters along a sequence of King models determined by changes of state resulting from collisions with the black holes. We include mass loss in collisions as well as the heating of the remaining bound stars, and determine the role which a finite number of stars plays in the variance of the energy input and mass loss. Several methods are used to determine the range of black hole masses and abundances excluded by survival of galactic globular clusters: simple order of magnitude estimates; collision-by-collision simulations of the energy input and mass loss of a stellar cluster; and a ‘smoothed’ Monte Carlo calculation of the evolution of cluster energy and mass. The results divide naturally into regimes of ‘small’ and ‘large’ black hole mass. ‘Small’ black holes do not destroy clusters in single collisions; their effect is primarily cumulative, leading to a relation between M_{bh} and f_{halo} , the fraction of the halo in black holes of mass M_{bh} , which is $f_{\text{halo}}M_{\text{bh}} < \text{constant}$ (up to logarithmic corrections). For $f_{\text{halo}} = 1$, we find $M_{\text{bh}} \lesssim 10^3 M_{\odot}$ by requiring survival of the same clusters studied by Moore (1993), who neglected cluster evolution, mass loss, and stochasticity of energy inputs in his estimates, but reached a similar conclusion. ‘Large’ black holes may not penetrate a cluster without disrupting it; their effect is mainly catastrophic (close collisions), but also partly cumulative (distant collisions). In the large M_{bh} limit, f_{halo} (but not M_{bh}) can be constrained by computing the probability that a cluster survives a combination of close, destructive encounters and distant, nondestructive encounters. We find that it is unlikely that $f_{\text{halo}} \gtrsim 0.3$ by requiring 50 per cent survival probability for Moore’s clusters over 10^{10} years.

Key words: Galaxy: general – globular clusters: general – Galaxy: halo – Galaxy: kinematics and dynamics – dark matter

1 INTRODUCTION

In this paper, we reexamine the idea that black holes compose a substantial fraction of the dark matter in the halo of our galaxy. Ostriker & Lacey (1985) originally proposed that heating by a significant population of halo black holes with masses $M_{\text{bh}} \sim 10^6 M_{\odot}$ could explain Wielen’s (1977) inference that the velocity dispersions $\sigma_{\star}$ of disk stars of age $t_{\star}$ follow the trend $\sigma_{\star} \propto t_{\star}^{1/2}$. Although subsequent work questioned the basis of this argument, both because it is unclear that $\sigma_{\star}$ rises as fast as $t_{\star}^{1/2}$ (e.g. Carlberg et al. 1985; Strömberg 1987; Gomez et al. 1990), and that black holes are needed to explain the observed trend of $\sigma_{\star}$ with $t_{\star}$ (e.g. Lacey 1991), the heating of disk stars remains a constraint on the masses and abundance of any hypothetical population of massive halo objects; in the notation of Wasserman & Salpeter (1994), if the halo consists of a fraction f_{halo} in the form of massive black holes, then the Ostriker & Lacey (1985) argument implies $f_{\text{halo}}M_{\text{bh}} \lesssim 10^{6.6} M_{\odot}$. Limits on the masses and abundances of massive black holes could have important cosmological implications (e.g. Loeb 1993; Umemura, Loeb, & Turner, 1993; Loeb & Rasio 1994).

Carr (1994) has reviewed various limits on baryonic or black hole dark matter in the halo of our Galaxy. (See also Carr & Sakellariadou 1998.) One of the most powerful constraints on black hole properties yet proposed was put forth by Moore (1993) who concluded that the survival of a set of relatively tenuous, low mass ($M \lesssim 4 \times 10^4 M_\odot$) globular clusters over a timespan $\gtrsim 7 \times 10^9$ years would be possible only if $M_{\text{bh}} \lesssim 10^3 M_\odot$. The argument employed by Moore (1993) was first applied to this problem by Wielen (1985), who explored perturbations of more massive clusters ($M \sim 10^6 M_\odot$) by heavier black holes, in the range advocated by Ostriker & Lacey (1985). [Klesson & Burkert (1995) extended Moore's and Wielen's calculations to a range of globular cluster properties.] The idea is to compute the heating of a globular cluster by passing black holes over a chosen timescale comparable to the cluster age [as noted above, Moore (1993) used 7×10^9 years, roughly half the age inferred for typical clusters]. If the total energy imparted by black hole perturbations is sufficiently large, then the cluster is said to be disrupted.

In view of the importance of this problem, we have begun a more comprehensive study of the disruption of the same set of globular clusters considered by Moore (1993) by a hypothetical population of halo black holes. This investigation aims to tighten up Moore's argument in several different ways. Some of the improvements are technical [e.g. Moore used a simple analytic form for the energy input to a cluster by a passing black hole which is only very approximate; Klesson & Burkert (1995) improved on his formula], but others are qualitative. Of particular importance are: (1) Moore computed the energy input for a 'static' cluster, whose structure was held fixed. Our calculations evolve the clusters along a King sequence. (2) Qualitatively, one expects a cluster that gains a large amount of energy compared to its initial binding energy from encounters with passing black holes to lose a large fraction of its mass, too. Our calculations include mass loss by the clusters. (3) Although one can get a rough idea of the survival probability by considering the mean heating of a cluster, the energy transfer process is actually stochastic, and the variance in the energy input may play an important role in final estimates of critical masses for disruption to be likely. (4) Moore's calculations pertain to black holes of relatively low mass, which are incapable of disrupting a cluster in a single perturbation. For black hole masses $M_{\text{bh}} \gtrsim MV_{\text{rel}}/\sigma_{\text{cl}}$, where σ_{cl} is the characteristic velocity dispersion of the cluster and V_{rel} its characteristic speed relative to the approaching black hole ($V_{\text{rel}} \sim 330 \text{ km s}^{-1}$ is a fair estimate), a single encounter at the cluster's tidal radius will likely destroy it. In this regime, one can obtain limits on the halo mass density in massive black holes, but not on M_{bh} (e.g. Wielen 1988; Wasserman & Salpeter 1994).

Our study combines analytic and Monte Carlo calculations. Our most complete results come from Monte Carlo simulations in which we simulate each encounter between a cluster and a black hole separately. For simplicity, we shall consider clusters at fixed galactocentric radius, as was done by Moore (1993). We model clusters using N point masses whose positions and velocities are chosen from a King distribution (see, e.g., Binney & Tremaine 1987). Velocity perturbations are computed for a given black hole impact parameter and speed relative to the cluster center of mass using the impulse approximation (e.g. Binney & Tremaine 1987). The change in velocity of the cluster center of mass is subtracted from each individual velocity perturbation to determine the change in cluster energy and mass as a consequence of the collision. Determining which stars are ejected is tricky in any scheme that does not employ a direct N body simulation of interparticle interactions, but as long as the perturbations in individual encounters are not excessive, it should be sufficient to designate for ejection those stars whose post-collision velocities exceed their local pre-collision escape speed. To find the new King model that describes the remaining cluster, we need its tidal radius in addition to its mass and energy after the black hole encounter; we get this by assuming that the tidal radius is proportional to $M^{1/3}$, consistent with our assumption of fixed galactocentric radius. Within these 'rules of the game' we simulate the evolution of the globular clusters studied by Moore (1993) over a timespan of 10^{10} years, or until they are disrupted, whichever comes first. Our simulations are more comprehensive than the Monte Carlo calculations of Klesson & Burkert (1995), who chose discrete black hole encounter times, relative velocities V_{rel} and impact parameters b from the appropriate probability distributions, but merely updated the cluster velocity dispersion by a completely deterministic amount $\Delta\sigma(b, V_{\text{rel}})$, without accounting for stochasticity of heating, mass loss or the change in internal cluster structure in individual encounters.

Several different criteria are employed to decide whether or not a cluster has been destroyed. Some of these may be called 'global' in that they depend on integrated properties of the cluster, such as total energy or total mass. We can regard a cluster as having been destroyed, for example, when its mass or energy or energy per mass has changed by a fractional amount in excess of some pre-set values (e.g. 0.5). Since we evolve models along a sequence of King models which is limited, clusters may also die when they reach the end of the sequence (see also Chernoff, Kochanek, & Shapiro 1986). The other criteria for cluster disruption to be used are 'local', in that they depend on changes in the properties of a cluster as a consequence of a single collision. Thus, if the mass or energy of a cluster changes by more than some pre-set fractional amounts in an encounter with a black hole, we shall regard it as disrupted. This should also allow us to control the inaccuracy of our criteria for determining the mass loss per encounter somewhat. We evaluate survival probabilities for the various criteria separately.

The Monte Carlo calculations outlined above allow us to study both the large and small M_{bh} regimes. In the large M_{bh} regime, where destruction occurs only after relatively few encounters, the Monte Carlo calculations ought to be reasonably fast computationally. However, for small M_{bh} where the encounters are more frequent (the regime focussed on by Moore 1993), we expect the Monte Carlo calculations to be more cumbersome, thus limiting the number of stars we can use in realizing clusters. Ideally, one would like to be able to represent the stars 'one-by-one' since the variances in energy input depend on

the number of cluster particles. Fortunately, the perturbations due to individual encounters are relatively gentle in the small M_{bh} regime, and the problem lends itself to a Fokker-Planck treatment, which promises to be faster (at the price – justified in our view – of losing the ability to resolve short timescale structure in the cluster evolution). We present a two dimensional Fokker-Planck scheme in which we follow cluster evolution in mass and energy.

The calculations reported here suffer from two or three principal deficiencies. Most important is their reliance on the sequence of King models and on the impulse approximation. In addition, one would like to interweave perturbations by a hypothetical population of black holes with well-established sources of heating, such as disk shocking (e.g. Ostriker, Spitzer, & Chevalier 1972; Spitzer & Chevalier 1973; Chernoff et al. 1986; Binney & Tremaine 1987); in the similar problem of wide binary evolution, the interplay of perturbations by stars, molecular clouds, and dark matter is known to be important (e.g. Retterer & King 1982; Bahcall, Hut, & Tremaine 1985; Weinberg, Shapiro, & Wasserman 1987; Wasserman & Weinberg 1991). Moreover, globular clusters evolve on their own as a consequence of internal relaxation, resulting in evaporation and energy changes even without external perturbations; limits on halo black hole properties may be altered when internally induced changes in state are accounted for properly. These important effects will be ignored here to concentrate solely on the cluster evolution due to the collisions with black holes. We shall study some of these issues in a subsequent paper (Murali et al. 1998).

2 QUALITATIVE OVERVIEW OF THE COLLISION PROCESS AND EVOLUTIONARY SCENARIOS

2.1 Approximations

Throughout this paper we employ the impulse approximation and ignore deflection of the perturbing black hole orbit from a straight line. For nonpenetrating encounters at impact parameter b and relative velocity V_{rel} , these approximations are valid when $\Omega b/V_{\text{rel}} \lesssim 1$. The characteristic frequency is roughly $\Omega \sim (GM/r_{\text{hm}}^3)^{1/2}$, where M is the cluster mass and r_{hm} is the cluster half mass radius, for the outer parts of the cluster where the tidal perturbation is strongest. If we set $V_{\text{rel}} = \xi_1 V_c$, where V_c is the galactic circular speed, and let $\sigma_0 = \xi_2 (GM/r_{\text{hm}})^{1/2}$ be the cluster's central one-dimensional velocity dispersion, then the impulse and straight line approximations should hold for $b/r_{\text{hm}} \lesssim (\xi_1 \xi_2) V_c / \sigma_0$. For the clusters studied in this paper, $V_c / \sigma_0 \gtrsim 10 - 100$, and the approximations only fail far outside r_{hm} .

For penetrating encounters, this assessment remains valid for ‘typical’ collisions, but may fail for perturbations of particles deep in the cluster core. This is because the characteristic frequency near the center of the cluster where the perturbations are largest is $\Omega \simeq (4\pi G \rho_0 / 3)^{1/2}$, where ρ_0 is the central mass density of the cluster. Adopting $b = r_{\text{hm}}$ as typical, we find that the impulse and straight line approximations hold for $V_{\text{rel}} \gtrsim 0.1 \times r_{\text{hm}}(\text{pc}) [\rho_0 / M_{\odot} \text{pc}^{-3}]^{1/2} \text{km s}^{-1}$. Although our approximations would fail for the most concentrated clusters, they remain true for the relatively tenuous ones studied here (and indeed for many ‘normal’ clusters).

We also ignore all processes influencing cluster evolution except perturbations by black holes even though our limits are based on survival probabilities over timescales long enough for tidal shocking and internal relaxation to be important. In more realistic simulations, including these effects could tighten limits on properties of hypothetical halo black holes.

2.2 Single Collisions

Consider a single collision between a globular cluster and a black hole which takes place in the halo of the Galaxy. Let the cluster have tidal radius r_t , N stars, King model parameter $W_0 \equiv \tilde{\psi}_0$, and total energy E . The black hole passes the cluster with a speed V_{rel} and an impact parameter b . Define the ‘collision parameter’, $\eta_c = (M_{\text{bh}}/M)(\sigma_0/V_{\text{rel}})$. In section §5.2, it will be shown that in the impulsive limit the energy input, mass loss and their variances are

$$\begin{aligned} \frac{\overline{\Delta E(b, V_{\text{rel}})}}{|E|} &= \eta_c^2 R_E(b, \eta_c, \tilde{\psi}_0) \\ \frac{\overline{\Delta M(b, V_{\text{rel}})}}{M} &= -\eta_c^2 R_M(b, \eta_c, \tilde{\psi}_0) \\ \frac{\sigma_{\Delta E}^2}{E^2} &= \frac{\eta_c^2}{N} R_{EE}(b, \eta_c, \tilde{\psi}_0) \\ \frac{\sigma_{\Delta M}^2}{M^2} &= \frac{\eta_c^2}{N} R_{MM}(b, \eta_c, \tilde{\psi}_0) \end{aligned} \quad (1)$$

where the b dependent radial functions have the approximate limits (see §5.3)

$$\begin{aligned} R_E &\simeq C_E(\tilde{\psi}_0) \left(\frac{r_t}{b} \right)^4 \\ R_M &\simeq C_M(\tilde{\psi}_0) \left(\frac{r_t}{b} \right)^4 \end{aligned}$$

$$\begin{aligned}
R_{EE} &\simeq C_{EE}(\tilde{\psi}_0) \left(\frac{r_t}{b}\right)^4 \\
R_{MM} &\simeq C_{MM}(\tilde{\psi}_0) \left(\frac{r_t}{b}\right)^4
\end{aligned} \tag{2}$$

when $b \gtrsim r_t$. As much of the mass in the cluster is contained within the core radius, one can actually use the b^{-4} dependence all the way into the core as a first approximation. Examples of the radial functions R_E , R_M , R_{EE} , and R_{MM} for impacts inside the cluster are given in §5.2.

For large enough M_{bh} , a single collision will suffice to disrupt the cluster. This will be referred to as the ‘high black hole mass limit’. Define ‘disruption’ to occur for an energy input of size $f|E|$. Then the cluster is destroyed in a single collision for

$$\frac{b}{r_t} \lesssim \frac{b_d}{r_t} \equiv \left(\frac{C_E}{f}\right)^{1/4} \eta_c^{1/2}. \tag{3}$$

A safe *overestimate* of the mass, M_{high} , at which the cluster may be disrupted is given by the black hole mass at which $b_d = r_t$. Using $V_{\text{rel}} = \xi_1 V_c$, $\xi_1 \simeq 1.5$, and the Galactic circular speed $V_c = 220 \text{ km s}^{-1}$, the result is

$$M_{\text{high}} = M \frac{V_c}{\sigma_0} \left(\frac{\xi_1^2 f}{C_E}\right)^{1/2} \tag{4}$$

which is bigger than M by a factor of $V_c/\sigma_0 \gg 1$.

The probability distribution for energy input (and mass loss) can be understood in the following qualitative terms. The ratio of the variance to the mean energy input is

$$\frac{\sigma_{\Delta E}^2}{(\Delta E)^2} = \frac{R_{EE}}{N \eta_c^2 R_E^2} \simeq \frac{C_{EE}}{N C_E^2} \frac{1}{\eta_c^2} \frac{b^4}{r_t^4}. \tag{5}$$

This becomes small for large N and small M_{bh} , but is also proportional to b^4 , insuring that $\sigma_{\Delta E} \geq \overline{\Delta E}$ for

$$\frac{b}{r_t} \gtrsim \frac{b_{\text{diff}}}{r_t} = \left(\frac{C_E^2}{C_{EE}}\right)^{1/4} N^{1/4} \eta_c^{1/2}. \tag{6}$$

Note that $b_{\text{diff}} > r_t$ for $M_{\text{bh}}/M \gtrsim V_{\text{rel}}/(\sqrt{N}\sigma_0)$. The distributions of mass loss and energy input are sharply defined about the mean for $b \lesssim b_{\text{diff}}$, and become ‘fuzzy’ for $b \gtrsim b_{\text{diff}}$. In particular, the energy input will *always* be sharply peaked about the mean at the destructive radius b_d , because $b_d/b_{\text{diff}} = (C_{EE}/f C_E N)^{1/4} \ll 1$. Hence the probability of getting an energy input which is *not* destructive when $b < b_d$ is quite small.

2.3 Evolution Over Many Collisions

If $M_{\text{bh}} \ll M_{\text{high}}$ (‘the small M_{bh} limit’), it will take many collisions to disrupt the cluster. Individual collisions only ‘tickle’ the cluster, and the evolution can be approximated by averaging over the effects of many collisions. In §5.4, we find that the mean rate of energy input, averaged over b and V_{rel} , takes the form

$$\frac{\langle \dot{E} \rangle}{|E|} = d_E(\eta_{c0}, \tilde{\psi}_0) \Gamma_0 \tag{7}$$

where $\Gamma_0 \equiv n_{\text{bh}} \pi r_t^2 V_c$, n_{bh} is the number density of black holes, $\eta_{c0} = (M_{\text{bh}}/M)(\sigma_0/V_c)$ and $d_E(\eta_{c0}, \tilde{\psi}_0) \simeq \kappa_E(\tilde{\psi}_0) \ln(1/\eta_{c0}) \eta_{c0}^2$. The mean time for disruption, which we define here to occur at $\Delta E = f|E|$, is

$$T_{\text{disrupt}} = \frac{f|E|}{\langle \dot{E} \rangle} = \frac{f}{\Gamma_0 \kappa_E \ln(1/\eta_{c0}) \eta_{c0}^2} \propto \frac{1}{M_{\text{bh}}} \tag{8}$$

so that for very small M_{bh} it takes a long time to disrupt the cluster. Given a value for T_{disrupt} there is a certain critical value of M_{bh} , called $M_{\text{bh,crit}}$, above which the cluster is disrupted. The variance of the averaged energy input in a time T takes on the form (see §5.4)

$$\frac{\sigma_{\Delta E}^2}{E^2} = \frac{d_{EE}}{N} \Gamma_0 T \tag{9}$$

where $d_{EE}(\eta_{c0}, \tilde{\psi}_0) \simeq \kappa_{EE}(\tilde{\psi}_0) \ln(1/\eta_{c0}) \eta_{c0}^2$.

Given an ensemble of clusters with initial energy E_0 , a time $T \gg 1/\Gamma_0$ later the cluster energies will be roughly $E(T) = E_0 + \langle \Delta E(T) \rangle \pm \sigma_E(T)$. For small times the variance will dominate the mean corresponding to a very broad spread in cluster states but for sufficiently long times the ensemble will peak sharply about the mean. We find that for times $T \simeq T_{\text{disrupt}}$

$$\frac{\sigma_{\Delta E}^2(T_{\text{disrupt}})}{(\langle \Delta E(T_{\text{disrupt}}) \rangle)^2} \simeq \frac{\kappa_{EE}}{N \kappa_E} \frac{|E|}{\langle \Delta E(T_{\text{disrupt}}) \rangle} = \frac{\kappa_{EE}}{N f \kappa_E} \propto \frac{1}{N} \ll 1 \tag{10}$$




Figure 1. Schematic of fraction of cluster destroyed as a function of M_{bh}

so that the range of final states is always sharply defined about the mean for times long enough to disrupt the cluster when $M_{\text{bh}} \ll M_{\text{high}}$. The probability of survival, P_s , is given (here) by the fraction of clusters with $\Delta E < f|E_0|$. This fraction will change from unity to a minimum value in a ‘transition region’ with size, $\delta M_{\text{bh,crit}}$, dictated by the ratio $\sigma_{\Delta E}(T_{\text{disrupt}})/\langle \Delta E(T_{\text{disrupt}}) \rangle$ so that $\delta M_{\text{bh,crit}}/M_{\text{bh,crit}} \sim N^{-1/2} \ll 1$.

When $M_{\text{bh}} \gtrsim M_{\text{high}}$, any collision inside the destructive radius b_d will destroy the cluster. In the tidal limit, the expected number of destructive encounters, N_d , in the time T is

$$N_d = \left(\frac{C_E}{f} \right)^{1/2} \frac{\rho_{\text{bh}}}{M} \pi r_t^2 \sigma_0 T \quad (11)$$

where $\rho_{\text{bh}} = M_{\text{bh}} n_{\text{bh}}$ is the mass density in black holes. For an ensemble of clusters with initial energy E_0 , the destructive $b < b_d$ collisions act as a ‘sink’ for clusters while the nondestructive $b > b_d$ encounters give rise to a slower, diffusive energy change. The probability of survival, P_s , (which takes on its minimum value in the $M_{\text{bh}} \gtrsim M_{\text{high}}$ limit) is the product of $\exp(-N_d)$, the probability that no single destructive encounters occur, and the probability that all the $b > b_d$ collisions combined give $\Delta E < f|E_0|$. Since N_d does not depend on M_{bh} , only ρ_{bh} (or equivalently f_{halo}) can be constrained in the $M_{\text{bh}} \gtrsim M_{\text{high}}$ limit, not the black hole mass. In addition, the cluster evolution is quite stochastic in this regime as it depends on whether the destructive collisions do or do not occur.

The model for cluster evolution described in this section is summarized in the (purely illustrative) diagram of Fig. 1. The fraction of clusters destroyed in time T , called $f_{\text{dest}} = 1 - P_s$, changes from 0 to ~ 0.8 in a region of width $\sim 2000 M_\odot$ centered on $M_{\text{bh,crit}} \sim 5000 M_\odot$. Note that f_{dest} in this example does not asymptote at one, but instead reaches $f_{\text{dest}} \sim 0.8$ corresponding to $N_d \sim 1$.

Table 1. Parameters of the Globular Clusters

Cluster Name	M ($M_\odot$)	r_t (pc)	r_{core} (pc)	R_g (kpc)
AM 4	700	11.7	3.7	30.0
Arp 2	18000	75.0	9.5	20.4
NGC 5053	37700	72.0	10.9	16.7
NGC 7492	10400	45.4	4.5	18.7
Pal 4	24900	92.3	15.4	96.0
Pal 5	13700	76.4	13.9	16.5
Pal 13	3000	30.0	3.0	25.7
Pal 14	10400	117.5	22.4	69.9
Pal 15	15000	41.9	10.5	30.0

3 THE CLUSTER MODEL AND EVOLUTION OF THE CLUSTERS

The nondimensional King models (e.g. Binney & Tremaine 1987) are uniquely determined by the normalized central potential $\tilde{\psi}_0 \equiv \Psi(r=0)/\sigma_1^2 \equiv \Psi_0/\sigma_1^2$, which is the parameter W_0 of King (1966). The distribution function is given by

$$f_{\text{king}}(\mathbf{r}, \mathbf{v}) d^3\mathbf{r} d^3\mathbf{v} = d^3\mathbf{r} d^3\mathbf{v} \frac{\rho_1}{(2\pi\sigma_1^2)^{3/2}} \left(e^{(v_{\text{esc}}^2 - v^2)/(2\sigma_1^2)} - 1 \right). \quad (12)$$

for stellar positions $\mathbf{r}$, velocities $\mathbf{v}$, and local escape speed $v_{\text{esc}}(r)$. The dimensional King models can be specified by three independent quantities such as E , M , and r_t . Define $\nu(\tilde{\psi}_0)$ by $E = \nu(\tilde{\psi}_0) (GM^2/r_t)$. For $0.0 < \tilde{\psi}_0 < 8.5$, or $-0.60 > \nu > -2.13$, $\nu(\tilde{\psi}_0)$ is single-valued so that the King model is known given E , M , and r_t . A physical reason for excluding large $\tilde{\psi}_0$ is that simulations have shown King models become susceptible to gravothermal instability at $\tilde{\psi}_0 \gtrsim 7.40$ (Wiyanto et al. 1985). The clusters we study in this paper are not core collapsed.

A further restriction on the clusters is that they be tidally limited by the galaxy. To include the time-dependent effect of tidal stripping due to the galaxy would require detailed restricted three-body simulations for a range of both globular cluster orbits and orbits of stars in the clusters. In this paper we consider a first approximation in which the the galactic tidal field provides a relationship between r_t and M , but does not contribute a time-dependent perturbing force. For circular orbits, if we define r_t as the distance from the cluster center to the Lagrange point of the cluster plus galaxy potential, we get

$$\frac{M}{3r_t^3} \simeq \frac{M_g}{R_g^3} \quad (13)$$

for a point mass galaxy with mass M_g and galactocentric radius R_g , and we adopt $M/r_t^3 = \text{constant}$ for our clusters as they evolve due to collisions with black holes.

A weakness of our paper is its dependence on the King model sequence. As we shall see in §8, for some clusters black hole collisions may force $\tilde{\psi}_0 \rightarrow 0$ after only modest energy input and mass loss, leading to very tight (but somewhat artificial) bounds on black hole properties. We shall rectify this deficiency in a subsequent paper where cluster structure is not restricted to the King sequence (Murali et al. 1998).

We shall use two different sets of globular clusters to determine a maximum allowed black hole mass, $M_{\text{bh,crit}}$. First we examine the set of loosely bound globular clusters found in Moore (1993) and listed in Table 1, but then we also investigate a larger cluster perhaps more representative of the initial cluster population.

4 THE BLACK HOLE MODEL

Black holes of mass M_{bh} are assumed to compose a spherical halo with an isotropic velocity distribution. A fraction f_{halo} of the total halo mass is presumed to be in the black holes. When no value of f_{halo} is explicitly stated, $f_{\text{halo}} = 1$ is assumed. We model the black hole population as a singular isothermal sphere with one-dimensional velocity dispersion σ_{bh} and mass density

$$\rho_{\text{bh}}(R_g) = f_{\text{halo}} \frac{\sigma_{\text{bh}}^2}{2\pi G R_g^2} \quad (14)$$

(e.g. Binney & Tremaine 1987, Ch.4) with $\sigma_{\text{bh}} = V_c/\sqrt{2} = 220/\sqrt{2} \text{ km s}^{-1} \simeq 156 \text{ km s}^{-1}$. The number density of black holes at R_g is

$$n_{\text{bh}}(R_g) = f_{\text{halo}} \frac{V_c^2}{4\pi G M_{\text{bh}} R_g^2}. \quad (15)$$

We ignore rotation of the black hole halo, so their velocity distribution is $f(V_{\text{bh}}) = (\pi V_c^2)^{-3/2} \exp(-V_{\text{bh}}^2/V_c^2)$. For computing the impulsive mass loss and energy input to the cluster due to the collision, we need the distribution of relative speeds, V_{rel} . Let V_{cl} be the cluster velocity, so $V_{\text{rel}} = V_{\text{bh}} - V_{\text{cl}}$. The distribution of relative speeds between the halo of black holes and the cluster becomes

$$f(V_{\text{rel}})dV_{\text{rel}} = \frac{1}{\sqrt{\pi}} \frac{V_c}{V_{\text{cl}}} \left\{ \exp\left(-\frac{(V_{\text{rel}} - V_{\text{cl}})^2}{V_c^2}\right) - \exp\left(-\frac{(V_{\text{rel}} + V_{\text{cl}})^2}{V_c^2}\right) \right\} \frac{V_{\text{rel}}dV_{\text{rel}}}{V_c^2}. \quad (16)$$

In this paper the clusters are on circular orbits with $V_{\text{cl}} = V_c$.

The mean relative speed given by this distribution is

$$\langle V_{\text{rel}} \rangle = V_c \left\{ \left(x + \frac{1}{2x} \right) \text{erf}(x) + \frac{1}{\sqrt{\pi}} \exp(-x^2) \right\} \quad (17)$$

where $x \equiv V_{\text{cl}}/V_c$ and $\text{erf}(x) = (2/\sqrt{\pi}) \int_0^x \exp(-t^2)dt$. As $x \rightarrow 0$, $\langle V_{\text{rel}} \rangle \rightarrow 2V_c/\sqrt{\pi}$ (the Gaussian result), as $x \rightarrow \infty$, $\langle V_{\text{rel}} \rangle \rightarrow V_{\text{cl}}$, and for $V_{\text{cl}} = V_c$ we find $\langle V_{\text{rel}}(x=1) \rangle/V_c \equiv \xi_1 \simeq 1.47$. Note that the distribution in eq. (16) is different from that used by Moore (1993) and Klesson & Burkert (1995). In addition, they chose to approximate the differential rate of collisions $d\Gamma = n_{\text{bh}} 2\pi b db f(V_{\text{rel}}) V_{\text{rel}} dV_{\text{rel}}$ by $n_{\text{bh}} 2\pi b db f(V_{\text{rel}}) V_c dV_{\text{rel}}$, which will lead to errors in the number of collisions and the rate of energy input and mass loss to the cluster.

For a globular cluster at R_g , there will be a certain value of M_{bh} below which there will be more than one black hole inside the cluster on average at any time (the ‘many-body’ limit). The number of black holes inside a cluster is $N_{\text{inside}} = n_{\text{bh}} 4\pi r_t^3/3 = (V_c^2 r_t^3)/(3GM_{\text{bh}} R_g^2)$. Since $r_t \sim GM/\sigma_0^2$, eq. 13 and 14 give $3(V_c^2/R_g^2)(r_t^2/\sigma_0^2) \sim 1$ and so $N_{\text{inside}} \simeq M/(10M_{\text{bh}})$. This exceeds one for $M_{\text{bh}} < 0.1M$ which we will see is close to black hole mass limits for some clusters. The mathematical description of the energy input is complicated in this regime since the duration of a collision $\sim r_{\text{hm}}/V_{\text{rel}}$ is *longer* than the time between collisions $\sim (n_{\text{bh}} \pi r_{\text{hm}}^2 V_{\text{rel}})^{-1}$.

5 THE IMPULSIVE ENERGY INPUT FOR A SINGLE ENCOUNTER

In the impulse and straight line approximations, the velocity change of a star due to the passage of a black hole is

$$\Delta \mathbf{v} = -\frac{2GM_{\text{bh}} \mathbf{s}}{V_{\text{rel}} s^2} \quad (18)$$

where $\mathbf{s} \perp \mathbf{V}_{\text{rel}}$ is the projected vector from the black hole to the star at closest approach. Since the stellar velocity $v \ll V_{\text{cl}}$ and $v \ll V_{\text{bh}}$, we may neglect v in the relative velocity so that $\mathbf{V}_{\text{rel}} \simeq \mathbf{V}_{\text{bh}} - \mathbf{V}_{\text{cl}}$. In this approximation, all stars receive a velocity kick in the same plane perpendicular to $\mathbf{V}_{\text{rel}}$. The density of cluster stars can be projected onto this plane; then for a star at projected position $\mathbf{R}$ relative to the cluster center, $\mathbf{s} = \mathbf{R} - \mathbf{b}$, where $\mathbf{b}$ is the impact parameter of the black hole relative to the cluster center.

The cluster is destroyed for impacts with $\Delta v \gtrsim \sigma_0$. An order of magnitude estimate of the ratio of these two speeds is

$$\frac{\Delta v}{\sigma_0} \sim \frac{M_{\text{bh}}}{M} \frac{\sigma_0}{V_{\text{rel}}} \frac{r_t^4}{b^4} \equiv \eta_c \frac{r_t^4}{b^4} \quad (19)$$

For a penetrating impact with $b \lesssim r_t$, the cluster will be destroyed if $\eta_c \gtrsim 1$. As the velocity kick, energy input, etc. in the impulsive limit must scale as $M_{\text{bh}}/V_{\text{rel}}$, η_c is a convenient dimensionless measure of the destructiveness of the collision.

Next let star i have a mass m_i and a velocity $\mathbf{v}_i$ with respect to the center of mass before the collision. The energy of the cluster before the collision is then

$$E_{\text{before}} = \sum_{i=1}^N m_i \left(\frac{1}{2} v_i^2 + \frac{1}{2} \phi_i \right) \quad (20)$$

where $\phi_i = -GM/r_i - \psi_i$ is the gravitational potential, and $v_{\text{esc},i} = \sqrt{2\psi_i}$ is the escape speed for star i . The velocity kick relative to the center of mass is

$$\delta \mathbf{v}_i = \Delta \mathbf{v}_i - \overline{\Delta \mathbf{v}} \equiv \Delta \mathbf{v}_i - \frac{1}{N} \sum_{i=1}^N \Delta \mathbf{v}_i \quad (21)$$

where $\overline{\Delta \mathbf{v}}$ is computed for a continuous cluster in Appendix A. Star i is ejected if $(\mathbf{v}_i + \delta \mathbf{v}_i)^2 \geq v_{\text{esc},i}^2$. The total amount of mass ejected from the cluster can then be formally written down as

$$\Delta M = - \sum_{i=1}^N m_i \theta_i(\text{ej}) \quad (22)$$

where $\theta_i(\text{ej}) \equiv \theta[(\mathbf{v}_i + \delta \mathbf{v}_i)^2 - v_{\text{esc},i}^2]$ is a useful shorthand notation; similarly

$\theta_i(\mathbf{b}) \equiv \theta[v_{\text{esc},i}^2 - (\mathbf{v}_i + \delta\mathbf{v}_i)^2]$ for the bound stars. We neglect the possibility that stars unbound by our criterion remain near the cluster for a long time, possibly to become bound once again in a subsequent encounter with a black hole [Spitzer (1987) discusses orbits of this type].

To find the change in cluster energy we need the kinetic and potential energy changes, ΔT and ΔV , respectively. The change in kinetic energy arises both from the kicks to the bound stars and the loss of the energy of the ejected stars:

$$\Delta T = -\frac{1}{2} \sum_{i=1}^N m_i v_i^2 \theta_i(\text{ej}) + \sum_{i=1}^N m_i \left(\mathbf{v}_i \cdot \delta\mathbf{v}_i + \frac{1}{2} \delta v_i^2 \right) \theta_i(\mathbf{b}). \quad (23)$$

The change in the potential energy is $\Delta V = V_{\text{after}} - V_{\text{before}}$, where the potential energy before the collision is

$$V_{\text{before}} = -\frac{G}{2} \sum_{i=1}^N \sum_{j \neq i=1}^N \frac{m_i m_j}{r_{ij}} [\theta_i(\mathbf{b}) \theta_j(\mathbf{b}) + 2\theta_i(\text{ej}) \theta_j(\mathbf{b}) + \theta_i(\text{ej}) \theta_j(\text{ej})]. \quad (24)$$

and potential energy of the bound stars remaining afterwards is

$$V_{\text{after}} = -\frac{G}{2} \sum_{i=1}^N \sum_{j \neq i=1}^N \theta_i(\mathbf{b}) \theta_j(\mathbf{b}) \frac{m_i m_j}{r_{ij}} \quad (25)$$

for separation vectors $\mathbf{r}_{ij}$. Consequently,

$$\Delta V = \sum_{i=1}^N \sum_{j \neq i=1}^N \theta_i(\text{ej}) \frac{G m_i m_j}{r_{ij}} - \frac{1}{2} \sum_{i=1}^N \sum_{j \neq i=1}^N \theta_i(\text{ej}) \theta_j(\text{ej}) \frac{G m_i m_j}{r_{ij}} \quad (26)$$

$$= -\sum_{i=1}^N \theta_i(\text{ej}) m_i \phi_i - \frac{1}{2} \sum_{i=1}^N \sum_{j \neq i=1}^N \theta_i(\text{ej}) \theta_j(\text{ej}) \frac{G m_i m_j}{r_{ij}} \quad (27)$$

where ϕ_i still denotes the potential of star i from *before* the collision. The first term in eq. (27) is just the pre-collision potential energy of the ejected stars, and is $\mathcal{O}(\Delta M/M)$, the fractional mass loss. The second term is $\mathcal{O}(\Delta M/M)^2$, and will be smaller than the first as long as $\Delta M/M$ is small. In our calculations we neglect the smaller (and difficult to compute) second term, and evaluate the first using the King model potential at the positions of the ejected stars. To protect against gross inaccuracies, we terminate our simulations if $|\Delta M/M| > 0.2$ in a single collision.

With these approximations, the energy change is

$$\Delta E = -\sum_{i=1}^N m_i \left(\frac{1}{2} v_i^2 + \phi_i \right) \theta_i(\text{ej}) + \sum_{i=1}^N m_i \left(\mathbf{v}_i \cdot \delta\mathbf{v}_i + \frac{1}{2} \delta v_i^2 \right) \theta_i(\mathbf{b}). \quad (28)$$

It should be noted that the energy change due to ejected stars is always positive, and hence mass loss always heats the cluster. In the sum over bound stars, the $\mathbf{v}_i \cdot \delta\mathbf{v}_i$ term can have either sign and hence can heat or cool. The mean of this term is nonzero but small since it depends on the mass loss to make the final distribution slightly anisotropic; it also contributes substantial variance. The $\frac{1}{2} \delta v_i^2$ term gives rise to the familiar mean heating (King 1966; Binney & Tremaine 1987).

5.1 The Model for Energy Input and Mass Loss

For the analytic estimates and Fokker-Planck calculations of cluster survival, we shall need formulae for the mean energy and mass changes, $\overline{\Delta E}$ and $\overline{\Delta M}$, their variances, and the cross-term $\overline{\Delta E \Delta M}$. To get them we use a model for energy input and mass loss which is due to Chernoff et al. (1986) (hereafter called C.K.S.). In this model, the portion of phase space from which mass escapes is identified. The energy and mass of the remaining cluster are then found as integrals over the distribution function of the remaining stars. This model relies only on the information given by the distribution function from *before* the collision; no detailed solution of the Vlasov equation is attempted. The higher order effects which arise from the detailed alteration of the distribution function by the perturbing black holes are estimated and define the error in our method.

We shall also extend the C.K.S. model slightly by considering the fluctuations in the number of stars lost in any collision resulting from a finite number N of stars. Before the collision, the phase space density of the cluster is given by the King model in eq. (12). After the collision, we do not have a full expression for the phase space density including the effect of the change in velocity to all the stars, but we do know that all parts of phase space for which $\theta(\text{ej}) = 1$ will have their mass ejected. Define $f_{<} = f_{\text{king}} \theta(\mathbf{b})$ and $f_{>} = f_{\text{king}} \theta(\text{ej})$ which are nonzero only for bound and unbound stars, respectively. The average of a quantity x over the bound, or ejected, stars will be written

$$\langle x \rangle_{<} = \frac{\int d^6 \Gamma f_{<} x}{\int d^6 \Gamma f_{<}}, \quad \langle x \rangle_{>} = \frac{\int d^6 \Gamma f_{>} x}{\int d^6 \Gamma f_{>}} \quad (29)$$

respectively. If a symbol $<$ or $>$ is not specified, it means that the quantity is averaged over the entire phase space.

The probability that a star is in the portion of phase space from which mass is ejected is $p_{\text{ej}} = \langle \theta(\text{ej}) \rangle$, which is an integral over the cluster model and is independent of N . We then suppose that the distribution for losing N_{ej} stars out of the total N is given by

$$P(N_{\text{ej}}) = \frac{(p_{\text{ej}}N)^{N_{\text{ej}}}}{N_{\text{ej}}!} \exp(-p_{\text{ej}}N). \quad (30)$$

In averaging the expressions of §5, we will only need the moments $\langle N_{\text{ej}} \rangle = Np_{\text{ej}}$ and $\langle N_{\text{ej}}^2 \rangle = (Np_{\text{ej}})^2 + Np_{\text{ej}}$.

5.2 Moments of ΔE and ΔM for Individual Collisions

In this section, we integrate the first and second moments of ΔE and ΔM from §5 over the King cluster model and the distribution for the number of ejected stars $P(N_{\text{ej}})$. These averages will be denoted by an overbar to distinguish them from the averages over bound and ejected stars defined in the previous section.

The mean mass loss and energy input are

$$\frac{\overline{\Delta M(b, \eta_c, \tilde{\psi}_0)}}{M} = -\langle \theta(\text{ej}) \rangle \quad (31)$$

and

$$\begin{aligned} \frac{\overline{\Delta E(b, \eta_c, \tilde{\psi}_0)}}{|E|} &= \frac{M}{|E|} \left\{ -p_{\text{ej}} \left\langle \frac{1}{2}v^2 + \phi \right\rangle_{>} + (1 - p_{\text{ej}}) \left\langle \mathbf{v} \cdot \delta \mathbf{v} + \frac{1}{2}\delta v^2 \right\rangle_{<} \right\} \\ &= \frac{M}{|E|} \left\{ -p_{\text{ej}} \left\langle \frac{1}{2}v^2 + \phi \right\rangle_{>} + \left\langle \mathbf{v} \cdot \delta \mathbf{v} + \frac{1}{2}\delta v^2 \right\rangle_{<} \right\} + \mathcal{O}(\eta_c^4). \end{aligned} \quad (32)$$

Consistent with our neglect of terms $\propto (\Delta M/M)^2$ (and smaller), we only retain contributions at $\mathcal{O}(\eta_c^2)$ and $\mathcal{O}(\eta_c^3)$; note that $p_{\text{ej}} \propto \eta_c^2$, $|\delta v| \propto \eta_c$, and consequently $\langle \mathbf{v} \cdot \delta \mathbf{v} \rangle_{<}$ – which would vanish in absence of mass loss – is $\propto \eta_c^3$. To the same accuracy, the variance of ΔM is

$$\frac{\sigma_{\Delta M}^2(b, \eta_c, \tilde{\psi}_0, N)}{M^2} = \frac{p_{\text{ej}}}{N} \quad (33)$$

and the variance of ΔE is

$$\begin{aligned} \frac{\sigma_{\Delta E}^2(b, \eta_c, \tilde{\psi}_0, N)}{E^2} &= \frac{M^2}{NE^2} \left[(1 - p_{\text{ej}}) \left\langle \left(\mathbf{v} \cdot \delta \mathbf{v} + \frac{1}{2}\delta v^2 \right)^2 \right\rangle_{<} + p_{\text{ej}} \left\langle \left(\frac{1}{2}v^2 + \phi \right)^2 \right\rangle_{>} \right. \\ &\quad \left. - \left(-p_{\text{ej}} \left\langle \frac{1}{2}v^2 + \phi \right\rangle_{>} + (1 - p_{\text{ej}}) \left\langle \mathbf{v} \cdot \delta \mathbf{v} + \frac{1}{2}\delta v^2 \right\rangle_{<} \right)^2 \right] \\ &\simeq \frac{M^2}{NE^2} \left[\langle (\mathbf{v} \cdot \delta \mathbf{v})^2 \rangle_{<} + p_{\text{ej}} \left\langle \left(\frac{1}{2}v^2 + \phi \right)^2 \right\rangle_{>} \right]. \end{aligned} \quad (34)$$


Finally, the mixed moment is

$$\begin{aligned} \frac{\sigma_{\Delta E \Delta M}^2(b, \eta_c, \tilde{\psi}_0, N)}{M|E|} &\equiv \frac{\overline{(\Delta M \Delta E)(\eta_c, \tilde{\psi}_0, N)}}{M|E|} - \frac{\overline{\Delta M}(\eta_c, \tilde{\psi}_0)}{M} \frac{\overline{\Delta E}(\eta_c, \tilde{\psi}_0)}{|E|} \\ &= \frac{M}{N|E|} \left[+p_{\text{ej}} \left\langle \frac{1}{2}v^2 + \phi \right\rangle_{>} \right], \end{aligned} \quad (35)$$

keeping only terms $\mathcal{O}(\eta_c^3)$ and larger.

We have calculated the integrals needed to evaluate these moments using Monte Carlo methods. The velocity kick is evaluated using eq. (18) and the center of mass velocity kick is given in eq. (A1). A sample of the results for $\tilde{\psi}_0 = 2.0, 4.0, 6.0$ and $\eta_c = 10^{-2}$ are plotted in Figs. 2, 3, and 4. For $\tilde{\psi}_0 = 2.0, 4.0$, and 6.0 , the core radii are at $r_{\text{core}}/r_t = 0.2, 0.1$, and 0.05 , respectively, while the half mass radii are at $r_{\text{hm}}/r_t = 0.3, 0.2$, and 0.15 , respectively. The Monte-Carlo integrals were computed to an estimated fractional error of 5 per cent.

The separate contributions to the energy input in eq. (32) have been plotted in Fig. 2. The contribution of mass loss is comparable to that due to the $\delta v^2/2$ heating usually considered. Note too that the $\langle \mathbf{v} \cdot \delta \mathbf{v} \rangle$ term is surprisingly large for small b encounters, contributing ~ 5 per cent to the energy input for impacts near the cluster center. The variance in eq. (34) is also broken up into the heating part, $\langle (\mathbf{v} \cdot \delta \mathbf{v})^2 \rangle_{<}$ and the mass loss part, $p_{\text{ej}} \left\langle \left(\frac{1}{2}v^2 + \phi \right)^2 \right\rangle_{>}$, which are comparable in size.

Figure 2. $\langle \Delta E(b, \eta_c) \rangle / |E|$.

5.3 The Tidal Limit


The calculation of $\overline{\Delta E}$, $\overline{\Delta M} = \sqrt{N} \sigma_{\Delta M}$, $\sigma_{\Delta E}^2$, and $\sigma_{\Delta E \Delta M}^2$ is simplified considerably in the tidal limit, partly because mass loss is restricted to particles with speeds very close to the escape speed. Spitzer (1958) computed $\overline{\Delta E}$ without mass loss, however, mass loss contributes significantly to $\overline{\Delta E}$, sometimes exceeding the Spitzer term. The expressions found below illustrate the importance of mass loss explicitly and are also useful for understanding disruption of clusters by high mass black holes.

Formally, we should expand in the two parameters η_c and $r_t/b \ll 1$. Here, we keep only the lowest powers of η_c and $\beta = r_t/b$. In this approximation, we find

$$\frac{\overline{|\Delta M|}(b, \eta_c, \tilde{\psi}_0)}{M} = -C_M \frac{\eta_c^2}{\beta^4} \quad (36)$$

where

$$C_M = \frac{8}{9\sqrt{\pi}} \frac{\mu_t}{\xi_t^4} \frac{\tilde{\psi}_0^{7/2}}{f_c g_c} \int_0^{\xi_t} d\xi \xi^4 \Theta(\xi)^{3/2}, \quad (37)$$

Figure 3. $|\langle \Delta M(b, \eta_c) \rangle|/M$.

and $\xi = r/a$, $\xi_t = r_t/a$, $a = (\Psi_0/4\pi G\rho_0)^{1/2}$, $\mu_t = M/M_0$, $M_0 = 4\pi\rho_0 a^3$, $f_c = \rho_0/\rho_1$, $g_c = \sigma_0^2/\sigma_1^2$, and $\Theta(\xi) = \Psi(\xi)/\Psi_0$ [ρ_1 and σ_1 are defined in eq. (12)]. Hence $\sigma_{\Delta M}^2/M^2 = p_{ej}/N = (C_{MM}/N)(\eta_c^2/\beta^4)$ where $C_{MM} = C_M$. To $\mathcal{O}(\eta_c^2)$,

$$\frac{\overline{\Delta E(b, \eta_c, \tilde{\psi}_0)}}{|E|} \simeq \frac{M}{|E|} \left\{ p_{ej} \frac{GM}{r_t} + \left\langle \frac{1}{2} \delta v^2 \right\rangle \right\} \simeq p_{ej} \frac{GM^2}{r_t |E|} + \frac{4}{3} \frac{G^2 M_{bh}^2 M \langle r^2 \rangle}{|E| V_{rel}^2 b^4} \equiv C_E \frac{\eta_c^2}{\beta^4} \quad (38)$$

with

$$C_E = C_M \frac{GM^2}{r_t |E|} + \frac{4}{3} \frac{G^2 M^3 \langle r^2 \rangle}{|E| \sigma_0^2 r_t^4}. \quad (39)$$

In addition to the Spitzer energy loss $\langle \delta v^2/2 \rangle$, C_E includes the energy carried away by the massloss, $\Delta M = -p_{ej}M$, from near the escape surface, where the energy per mass is $-GM/r_t$. Note that the ejected mass can come from any spatial position in the cluster as long as the velocity is close enough to the escape velocity. Indeed, the mean radius from which mass is lost




Figure 4. $N\sigma_{\Delta E}^2(b, \eta_c)/E^2$.

depends weakly on cluster concentration, and is roughly $2r_t/3$ for $0 \leq \tilde{\psi}_0 \leq 8.5$. The variance to the energy input is

$$\frac{\sigma_{\Delta E}^2(b, \eta_c, \tilde{\psi}_0, N)}{E^2} = \frac{1}{N} p_{\text{ej}} \left(\frac{GM^2}{|E|r_t} \right)^2 + \frac{8}{9} \frac{1}{N} \frac{G^2 M_{\text{bh}}^2 M^2}{E^2 V_{\text{rel}}^2 b^4} \langle r^2 v^2 \rangle \equiv \frac{C_{EE}}{N} \frac{\eta_c^2}{\beta^4} \quad (40)$$

with

$$C_{EE} = C_M \left(\frac{GM^2}{|E|r_t} \right)^2 + \frac{8}{9} \frac{G^2 M^4}{E^2 \sigma_0^2 r_t^4} \langle r^2 v^2 \rangle, \quad (41)$$

and the mixed moment is

$$\frac{\sigma_{\Delta E \Delta M}^2(b, \eta_c, \tilde{\psi}_0, N)}{M|E|} = \frac{M}{N|E|} \left[+p_{\text{ej}} \left\langle \frac{1}{2} v^2 + \phi \right\rangle \right] \simeq -\frac{M}{N|E|} p_{\text{ej}} \frac{GM}{r_t} \equiv -\frac{C_{EM}}{N} \frac{\eta_c^2}{\beta^4} \quad (42)$$

with $C_{EM} = C_M(GM^2)/(|E|r_t)$.

Another extremely important quantity will be the change in $\nu \equiv (Er_t)/(GM^2)$ in the tidal limit. Using the fact that the cluster is tidally limited, ν can be expressed as $\nu = (r_{t,0}/(GM_0^{1/3})) \times (E/M^{5/3})$ where the subscript refers to some reference




Figure 5. Dimensionless coefficients for fractional energy input and mass loss in the tidal limit.

value. The change in ν is then

$$\Delta \nu = \nu \left(\frac{\Delta E}{E} - \frac{5}{3} \frac{\Delta M}{M} \right) = |\nu| \left(C_E - \frac{5}{3} C_M \right) \eta_c^2 \frac{r_t^4}{b^4} \quad (43)$$

which can be written in the usual form

$$\frac{\overline{\Delta \nu(b, \eta_c, \tilde{\psi}_0)}}{|\nu|} = C_\nu \frac{\eta_c^2}{\beta^4} \quad (44)$$

with $C_\nu = C_E - 5C_M/3$. The variance of the mean change in ν is given by

$$\frac{\sigma_{\Delta \nu}^2(b, \eta_c, \tilde{\psi}_0, N)}{\nu^2} = \frac{\sigma_{\Delta E}^2}{E^2} + \frac{5}{3} \frac{\sigma_{\Delta E \Delta M}^2}{M|E|} + \frac{25}{9} \frac{\sigma_{\Delta M}^2}{M^2} = \frac{C_{\nu\nu}}{N} \frac{\eta_c^2}{\beta^4} \quad (45)$$

where $C_{\nu\nu} \equiv C_{EE} - 5C_{EM}/3 + 25C_{MM}/9$.

The seven coefficients for the tidal limit are plotted in Figs. 5 and 6 as a function of $\tilde{\psi}_0$. All the curves show a monotonic decrease as $\tilde{\psi}_0$ increases over the range $\tilde{\psi}_0 \in (0.0, 8.5)$. Most noticeable is that C_ν becomes less than zero at roughly $\tilde{\psi}_0 \simeq 5.5$. Since $\nu = \nu(\tilde{\psi}_0)$, whether an impact will make the cluster more or less concentrated is determined by the value of $\tilde{\psi}_0$ for that cluster. Clusters with $\tilde{\psi}_0 < 5.5$ are driven toward dissolution and clusters with $\tilde{\psi}_0 > 5.5$ are driven toward core collapse. This agrees well with CKS who derived the same result in the context of perturbations by giant molecular clouds.

For penetrating encounters, the situation is quite different. Over almost the entire range of $\tilde{\psi}_0$ and η_{c0} , the diffusion coefficients yield positive changes in ν , implying dissolution.

5.4 Evaluation of the Diffusion Coefficients

In §5.2, we found $\overline{\Delta E}$, $\overline{\Delta M}$, $\sigma_{\Delta E}^2$, $\sigma_{\Delta M}^2$, and $\sigma_{\Delta E \Delta M}^2$ for individual collisions as a function of impact parameter b and relative velocity V_{rel} . When M_{bh} is small enough that the changes in the cluster properties are always small for single impacts, then we may average over many encounters to find the changes in E and M over a time period which includes many collisions, but for which the changes in cluster properties are still small.




Figure 6. Dimensionless coefficients for fractional changes in ν in the tidal limit.

We weight the formulae for the various moments by the differential rate of collisions

$$d\Gamma = n_{\text{bh}} 2\pi b db f(V_{\text{rel}}) V_{\text{rel}} dV_{\text{rel}} \quad (46)$$

and integrate over b and V_{rel} where $f(V_{\text{rel}})$ is given in eq. (16). The results may be written

$$\begin{aligned} D_E &= \int d\Gamma \overline{\Delta E} = \Gamma_0 |E| d_E \\ D_M &= \int d\Gamma \overline{\Delta M} = -\Gamma_0 M d_M \\ D_{EE} &= \int d\Gamma \sigma_{\Delta E}^2 = \frac{1}{N} \Gamma_0 E^2 d_{EE} \\ D_{MM} &= \int d\Gamma \sigma_{\Delta M}^2 = \frac{1}{N} \Gamma_0 M^2 d_M \\ D_{EM} &= \int d\Gamma \sigma_{\Delta E \Delta M}^2 = -\frac{1}{N} \Gamma_0 |E| M d_{EM} \end{aligned} \quad (47)$$

where $\Gamma_0 = n_{\text{bh}} \pi r_t^2 V_c$, and d_E , d_M , d_{EE} , and d_{EM} are dimensionless functions of $\tilde{\psi}_0$ and $\eta_{c0} = (M_{\text{bh}}/M)(\sigma_0/V_c)$. Note that minus signs have been inserted in D_M and D_{EM} to make d_M and d_{EM} positive.

The diffusion coefficients d_E , d_M , d_{EE} , and d_{EM} involve eight-dimensional integrals over phase space, b , and V_{rel} . Integration over two velocity angles is elementary, and the integral over velocity magnitude was done using Runge-Kutta. The remaining five integrals (three spatial coordinates, b , and V_{rel}) were done via Monte Carlo (to an accuracy of 5 per cent). The value of b_{max} was taken to be $10r_t$.

For use in our Fokker-Planck calculations, the four diffusion coefficients were tabulated for almost all the values in the range $\tilde{\psi}_0 = 0.0 \rightarrow 8.5$ and $\eta_{c0} = 0.0001 \rightarrow 0.5$. Due to the extremely long integration times for small values of η_{c0} and large $\tilde{\psi}_0$, a few values were obtained using analytic fitting formulae of the form $d_E = \kappa_E \ln(1/\eta_{c0}) \eta_{c0}^2$, $d_M = \kappa_M \ln(1/\eta_{c0}) \eta_{c0}^2$, $d_{EE} = \kappa_{EE} \ln(1/\eta_{c0}) \eta_{c0}^2$, and $d_{EM} = \kappa_{EM} \ln(1/\eta_{c0}) \eta_{c0}^2$, which represented the data well; the coefficients for these fits are given in Table 2. The fits were generally good to $\lesssim 5-30$ percent and are useful for quick estimates, although for more precise

Table 2. Fitting Formula Coefficients for d_E , d_M , d_{EE} , and d_{EM} .

$\tilde{\psi}_0$	κ_E	$\hat{\kappa}_E$	κ_M	$\hat{\kappa}_M$	κ_{EE}	$\hat{\kappa}_{EE}$	κ_{EM}	$\hat{\kappa}_{EM}$
0.0	12.8	70.5	4.14	22.8	20.8	114	7.29	40.1
0.5	13.4	80.5	4.34	26.1	23.3	140	7.90	47.5
1.0	11.0	73.2	3.38	22.5	18.3	122	5.89	39.2
1.5	10.1	75.3	3.10	23.1	16.8	126	5.31	39.6
2.0	7.05	60.1	2.06	17.6	10.0	85.3	3.07	26.2
2.5	7.68	76.2	2.24	22.3	12.2	122	3.56	35.4
3.0	6.42	75.8	1.85	21.9	9.64	114	2.72	32.2
3.5	5.42	78.2	1.51	21.8	7.95	115	2.09	30.1
4.0	4.85	88.0	1.39	25.2	7.33	133	1.93	35.0
4.5	3.80	89.5	1.06	24.9	5.55	131	1.33	31.3
5.0	2.85	90.2	0.740	23.4	4.05	128	0.985	31.2
5.5	2.33	102	0.702	30.8	3.36	147	0.780	34.1
6.0	1.88	117	0.603	37.7	2.73	171	0.631	39.4
6.5	1.40	126	0.499	45.0	2.03	183	0.489	44.0
7.0	1.06	137	0.379	48.8	1.56	201	0.329	42.4
7.5	0.836	141	0.363	61.4	1.24	209	0.308	52.0
8.0	0.677	134	0.320	63.2	0.971	192	0.243	48.0
8.5	0.647	126	0.362	70.7	0.928	181	0.277	54.0

Note: The dimensionless diffusion coefficients are approximated by $d = \kappa \ln(1/\eta_{c0}) \eta_{c0}^2$ and $\hat{d} = \hat{\kappa} \ln(1/\hat{\eta}_{c0}) \hat{\eta}_{c0}^2$ where $\hat{d} = d(r_t^2/r_{\text{hm}}^2)$ and $\hat{\eta}_{c0} = \eta_{c0}(GM/r_{\text{hm}}\sigma_0^2)^{1/2}$.

numerical work interpolation on tabulated values was used. The values generated using the fitting formulae were used only for the very smallest values of η_{c0} for the cluster PAL 5, where the fitting formulae are most accurate. Fig. 7 displays the various coefficients for several values of η_{c0} which are shown increasing from bottom to top. The triangles represent data which was generated by the Monte Carlo program itself, while the open circles represent the data estimated using the fitting formula.

It can be seen that for a fixed value of η_{c0} , the diffusion coefficients in Fig. 7 vary by about an order of magnitude over the range $\tilde{\psi}_0 = 0.0 \rightarrow 8.5$. The coefficients for the fitting formulae show a similar range of variation. This spread can be reduced considerably if we recall that many quantities vary little over the King sequence when expressed in terms of the half-mass radius, r_{hm} . Choose a new unit of rate $\hat{\Gamma}_0 = n_{\text{bh}} \pi r_{\text{hm}}^2 V_c$ and a new collision parameter $\hat{\eta}_{c0} = \eta_{c0}(GM/r_{\text{hm}}\sigma_0^2)^{1/2}$. The constants $\hat{\kappa}$ in the new dimensionless diffusion coefficients $\hat{d} = \hat{\kappa} \ln(1/\hat{\eta}_{c0}) \hat{\eta}_{c0}^2 = d(r_t^2/r_{\text{hm}}^2)$ then vary by about a factor of two over the range $\tilde{\psi}_0 = 0.0 \rightarrow 8.5$, as can be seen in Table 2.

6 ‘SLOW HEATING’ LIFETIMES

6.1 The Fixed Cluster Approximation

When $M_{\text{bh}} \ll M_{\text{high}}$, the properties of a given cluster only change slightly over many encounters with black holes. In this limit, cluster survival implies an upper bound to $f_{\text{halo}} M_{\text{bh}}$ (up to logarithmic corrections), or M_{bh} assuming a certain value of f_{halo} . Very rough estimates of the critical black hole mass for survival over time T can be made if we neglect the evolution of the cluster profile over time T , and require the mean energy or mass change not to exceed some threshold over that time span (e.g. Moore 1993). The values found here are intended as benchmarks for comparison to the more detailed results, obtained in the next section, that include cluster evolution and statistical spreads in the energy and mass changes in time T .

For a fixed cluster profile, dE/dt , dM/dt , and $d\nu/dt$ are independent of time, and $\langle \Delta E(T) \rangle = \Gamma_0 |E| d_E T$ and $\langle \Delta M(T) \rangle = -\Gamma_0 M d_M T$, with the coefficients d_E and d_M defined and computed in §5.4. Choosing a time of $T = 10^{10}$ years as the time over which the clusters have been subjected to collisions, we defined three distinct values of $M_{\text{bh,crit}}$ by $\langle \Delta E(T, M_{\text{bh,crit}}) \rangle = |E|/2$, $\langle \Delta M(T, M_{\text{bh,crit}}) \rangle = -M/2$, and $\langle \Delta \nu(T, M_{\text{bh,crit}}) \rangle = [D_E/|E| - (5/3)(D_M/M)] |\nu| T = -0.6 - \nu$. The results are presented in Table 3 for the sample of nine weakly bound clusters employed by Moore (1993). Moore’s results correspond most closely to $M_{\text{bh,crit}}(E)$ as he used the criterion $\langle \Delta E(T = 7 \times 10^9 \text{ years}, M_{\text{bh,crit}}) \rangle = |E|$. When comparing to Moore’s results one should multiply $M_{\text{bh,crit}}$ by a factor of 2/0.7 to account for the larger value of T and smaller ΔE used here. Except for PAL 13, our results agree with Moore’s to within a factor of order a few, which is reasonably close considering the improvements made here (e.g. inclusion of mass loss, correct V_{rel} distribution).




Figure 7. Dimensionless diffusion coefficients. The triangles represent Monte Carlo results. The circles represent the incomplete data given by the fitting formulae. The values of η_{c0} shown are 0.0001, 0.00025, 0.0005, 0.00075, 0.001, 0.0025, 0.005, 0.0075, 0.01, 0.025, 0.05, 0.075, 0.1, 0.25, 0.5, increasing from bottom to top.

Table 3. Cluster Destruction for a Fixed Profile

Cluster Name	$M_{\text{bh,crit}}(E)/M_{\odot}$	$M_{\text{bh,crit}}(M)/M_{\odot}$	$M_{\text{bh,crit}}(\nu)/M_{\odot}$
AM 4	1500	7000	710
ARP 2	3100	13,000	4900
NGC 5053	4100	18,000	5200
NGC 7492	3400	15,000	6500
PAL 4	170,000	8,500,000	350,000
PAL 5	1100	4300	1100
PAL 13	3100	13,000	6300
PAL 14	14,000	68,000	15,000
PAL 15	9100	39,000	6100

Note: $M_{\text{bh,crit}}(E)/M_{\odot}$ is the critical black hole mass for the criterion $\delta E/|E| = 0.5$. $M_{\text{bh,crit}}(M)/M_{\odot}$ is the critical black hole mass for the criterion $|\delta M|/M = 0.5$. $M_{\text{bh,crit}}(\nu)/M_{\odot}$ is critical black hole mass for ν to go from its initial value to $\nu = -0.6$.

6.2 Gaussian Model with Evolution

In §6.1, two approximations were made. First, the cluster was treated as having a fixed profile for which the diffusion coefficients did not change over time. The second approximation was that the energy input and mass loss had sharply defined values over any time interval. In this section, the evolution of the cluster is followed over appropriately chosen intervals δt and a Gaussian distribution of energy input and mass loss is assumed.

The characteristic time for $N_{\min}$ collisions to occur inside $b_{\max}$ is

$$T_{\text{coll}} = \frac{N_{\min}}{n_{\text{bh}} \pi b_{\max}^2 \xi_1 V_c} \propto M_{\text{bh}} R_g^2 N_{\min}. \quad (48)$$

As most of the energy input is given by the penetrating encounters, a conservative estimate is to set $b_{\max} = r_t$ in T_{coll} . If $\epsilon_{\max}$ is the largest change allowed for the cluster in a time period, then a time T_{change} over which the cluster evolves significantly is given by (§5.4)

$$T_{\text{change}} = \frac{\epsilon_{\max}}{n_{\text{bh}} \pi r_t^2 V_c d_E} \propto \frac{\epsilon_{\max} R_g^2}{M_{\text{bh}}}. \quad (49)$$

The Gaussian approach is justified if there exists a δt such that $T_{\text{coll}} < \delta t < \delta T_{\text{change}}$, so the cluster properties change little over δt but enough collisions occur that the distribution of energy inputs and mass loss is approximately Gaussian. An estimate of the largest black hole mass, $M_{\text{bh,fp}}$, for which the Gaussian approach is justified is

$$M_{\text{bh,fp}} \lesssim M \frac{V_c}{\sigma_0} \left(\frac{\epsilon_{\max}}{N_{\min}} \right)^{1/2}, \quad (50)$$

or $\eta_{c0} \lesssim (\epsilon_{\max}/N_{\min})^{1/2}$. The critical mass can only be found in the present ‘diffusion’ approximation when $\epsilon_{\max} \ll 1$ and $N_{\min} \gtrsim 1$ so that the cluster must be destroyed for $\eta_{c0} \ll 1$.

Let the vector $\boldsymbol{\tau} = (\tau_1, \tau_2) = (\Delta E, \Delta M)$ where ΔE and ΔM are the energy input and mass loss over the time interval δt ; the expected values of ΔE and ΔM over the time δt are $\langle \tau_1 \rangle = \langle \Delta E(\delta t) \rangle$ and $\langle \tau_2 \rangle = \langle \Delta M(\delta t) \rangle$. The normalized distribution of energy input and mass loss is then

$$P(\boldsymbol{\tau}) d^2 \boldsymbol{\tau} = \frac{\sqrt{\det(\boldsymbol{\Xi})}}{2\pi} d^2 \boldsymbol{\tau} \exp \left\{ -\frac{1}{2} (\boldsymbol{\tau} - \langle \boldsymbol{\tau} \rangle)^T \boldsymbol{\Xi} (\boldsymbol{\tau} - \langle \boldsymbol{\tau} \rangle) \right\}, \quad (51)$$

where $\boldsymbol{\Xi}(\delta t)$ is the covariance matrix. The matrix $\boldsymbol{\Xi}^{-1}$ may be shown to be

$$\boldsymbol{\Xi}^{-1} = \begin{pmatrix} \langle (\tau_1 - \langle \tau_1 \rangle)^2 \rangle & \langle (\tau_1 - \langle \tau_1 \rangle)(\tau_2 - \langle \tau_2 \rangle) \rangle \\ \langle (\tau_1 - \langle \tau_1 \rangle)(\tau_2 - \langle \tau_2 \rangle) \rangle & \langle (\tau_2 - \langle \tau_2 \rangle)^2 \rangle \end{pmatrix};$$

inverting implies

$$\boldsymbol{\Xi} = \frac{1}{\det(\boldsymbol{\Xi}^{-1})} \begin{pmatrix} \langle (\tau_2 - \langle \tau_2 \rangle)^2 \rangle & -\langle (\tau_1 - \langle \tau_1 \rangle)(\tau_2 - \langle \tau_2 \rangle) \rangle \\ -\langle (\tau_1 - \langle \tau_1 \rangle)(\tau_2 - \langle \tau_2 \rangle) \rangle & \langle (\tau_1 - \langle \tau_1 \rangle)^2 \rangle \end{pmatrix}.$$

These expressions reduce to the standard sum of Gaussian terms when the correlation $\langle (\tau_1 - \langle \tau_1 \rangle)(\tau_2 - \langle \tau_2 \rangle) \rangle$ is zero. The eigenvalues of $\boldsymbol{\Xi}$ are $\lambda_{\pm} = \frac{1}{2} \left[\text{Tr}(\boldsymbol{\Xi}) \pm \sqrt{(\text{Tr}(\boldsymbol{\Xi}))^2 - 4 \det(\boldsymbol{\Xi})} \right]$.

In order to choose values for ΔE and ΔM , we define a new set of variables $\boldsymbol{\zeta} = (\zeta_+, \zeta_-)$ along the eigenvectors of $\boldsymbol{\Xi}$; the distribution of $\boldsymbol{\zeta}$ is

$$P(\boldsymbol{\zeta}) d^2 \boldsymbol{\zeta} = \frac{\sqrt{\lambda_+ \lambda_-}}{2\pi} d^2 \boldsymbol{\zeta} \exp \left\{ -\lambda_+ \zeta_+^2 / 2 - \lambda_- \zeta_-^2 / 2 \right\}, \quad (52)$$

and $\boldsymbol{\tau} - \langle \boldsymbol{\tau} \rangle = \boldsymbol{\Lambda} \boldsymbol{\zeta}$, where the unitary matrix

$$\boldsymbol{\Lambda} = \begin{pmatrix} \Lambda_{1+} & \Lambda_{1-} \\ \Lambda_{2+} & \Lambda_{2-} \end{pmatrix} = \frac{1}{\sqrt{(\lambda_- - \Xi_{22})^2 + \Xi_{12}^2}} \begin{pmatrix} \Xi_{12} & \lambda_- - \Xi_{22} \\ -(\lambda_- - \Xi_{22}) & \Xi_{12} \end{pmatrix}.$$

After choosing $\zeta_{\pm}$ from this Gaussian distribution, we multiply by $\boldsymbol{\Lambda}$ to get

$$\begin{aligned} \Delta E &= \langle \Delta E \rangle + \Lambda_{1+} \zeta_+ + \Lambda_{1-} \zeta_- \\ \Delta M &= \langle \Delta M \rangle + \Lambda_{2+} \zeta_+ + \Lambda_{2-} \zeta_-. \end{aligned} \quad (53)$$

In the limit of infinite numbers of collisions N_c in time δt (the limit in which the Gaussian model is equivalent to a 2-D Fokker-Planck equation)

$$\begin{aligned} \langle \tau_1 \rangle &= \Gamma_0 |E| d_E \delta t \\ \langle \tau_2 \rangle &= -\Gamma_0 M d_M \delta t \\ \Xi_{11}^{-1}(\delta t) &= \frac{1}{N} \Gamma_0 E^2 d_{EE} \delta t \\ \Xi_{12}^{-1}(\delta t) &= -\frac{1}{N} \Gamma_0 |E| M d_{EM} \delta t \\ \Xi_{22}^{-1}(\delta t) &= \frac{1}{N} \Gamma_0 M^2 d_{MM} \delta t. \end{aligned} \quad (54)$$

For finite N_c , these expressions are accurate to $\mathcal{O}(N_c^{-1})$.

Let us ignore the correlations of the change in mass and energy and focus solely on changes in cluster energy in order to get a qualitative feel for the ‘distribution of cluster states’. The Fokker-Planck equation for the distribution of energies, $P(E, t)$, of an ensemble of clusters is

$$\frac{\partial P}{\partial t} = -\frac{\partial}{\partial E}(D_E P) + \frac{1}{2} \frac{\partial^2}{\partial E^2}(D_{EE} P). \quad (55)$$

If we ignore changes in the cluster properties and take D_E and D_{EE} to be constant, this equation has the solution

$$P(E, t) = \frac{1}{(2\pi D_{EE} t)^{1/2}} \exp\left[-\frac{(E - E_0 - D_E t)^2}{2 D_{EE} t}\right] \quad (56)$$

where we have chosen the initial condition $E = E_0$ at $t = 0$. For small times $t < D_{EE}/D_E$, the width of the Gaussian is much larger than the mean implying a very spread out set of cluster states, and visa versa for times $t > D_{EE}/D_E$. As we have already shown in §2.3, for times $t \simeq T_{\text{disrupt}}$ the set of final states is sharply defined about the mean. Using the survival criterion $\Delta E < f|E_0|$, we find that the probability of survival after time t is given by

$$P_s = \int_{-\infty}^{E_0 + f|E_0|} dE P(E, t) = \frac{1}{2} \left[1 + \operatorname{erf}\left(\frac{f|E_0| - D_E t}{\sqrt{2 D_{EE} t}}\right) \right]. \quad (57)$$

The same equation would have resulted if we had used the two-dimensional Fokker-Planck equation for E and M . In terms of the fitting formulas of §5.4, the characteristic mass range, $\delta M_{\text{bh,crit}}$ over which P_s changes from one to zero is

$$\frac{\delta M_{\text{bh,crit}}}{M_{\text{bh,crit}}} \simeq \sqrt{\frac{\kappa_{EE}}{N f \kappa_E}} \ll 1 \quad (58)$$

so that in the $M_{\text{bh}} \ll M_{\text{high}}$ limit the value of $M_{\text{bh,crit}}$ found from just *one* history is likely to be close to the statistical mean.

6.3 Simulations and Results

We chose the accuracy parameters from the previous section to be $\epsilon_{\text{max}} = 0.01$ and $N_{\text{min}} = 10$. The time step was chosen so that the estimated change in energy would be 0.01. As the diffusion coefficients tended to increase as $\tilde{\psi}_0 \rightarrow 0$ and M decreased, the time step was decreased if two events in a row occurred with fractional changes in ν , M , or E greater than 0.01. Also, the time step was increased if the estimated number of events in δt was less than N_{min} . If $M_{\text{bh}} > M_{\text{bh,fp}}$, as described in the previous section, the simulation was stopped because the approximations had broken down.







For each of the nine weakly bound clusters from Moore (1993), values of M_{bh} were chosen and the cluster was evolved to one of the following outcomes: (1) the Fokker-Planck assumption broke down; (2) $T = 10^{10}$ years was reached and the cluster survived; (3) the quantity ν went out of the range $(-2.13, -0.6)$, signalling either core collapse or ‘dissolution’. Survival probabilities and their uncertainties were derived from the number of surviving clusters out of 1000 simulations using Bayesian arguments. The results are shown in Fig. 8 for Moore’s clusters as well as one more ‘normal’ cluster. Note that all curves asymptote to $f_{\text{dest}} = 1.0$ as $M_{\text{bh}} \rightarrow \infty$ because single-event destruction has been ignored in these simulations.

PAL 4 did *not* satisfy the Fokker-Planck assumption at the black hole mass required to disrupt it in $T = 10^{10}$ years. Since PAL 4 lies at such a great galactocentric distance, encounters are so infrequent that disruption in 10^{10} years requires individually destructive collisions. Hence a critical black hole mass $M_{\text{bh,crit}}$ cannot be determined by the slow heating approximation for PAL 4; the more detailed collision-by-collision encounter history of §8 is warranted for this case.

As for the rest of Moore’s clusters, we see that their values of $M_{\text{bh,crit}}$ agree well with the last column of Table 3; the values obtained from the Gaussian Monte-Carlo simulation are lower by less than a factor of two. As clusters are heated and lose mass, diffusion coefficients increase as $\tilde{\psi}_0 \rightarrow 0$, and η_{c0} decreases leading to smaller $M_{\text{bh,crit}}$ than for fixed cluster structure and mass. Also note that the fractional width, $\delta M_{\text{bh,crit}}/M_{\text{bh,crit}}$, of the ‘transition region’ over which $f_{\text{dest}} = 0.1 \rightarrow 0.9$ varies as $N^{-1/2}$, as predicted in §2.3.

We also simulated a large cluster with $M = 10^6 M_{\odot}$, $N = 1.4 \times 10^6$, $r_t = 50$ pc, $r_{\text{core}} = 5$ pc, and $R_g = 10$ kpc; hence $\tilde{\psi}_0 = 4.85$ and the central one-dimensional velocity dispersion $\sigma_0 = 9.64$ km s $^{-1}$. Fig. 8 shows P_s as a function of M_{bh} for this cluster. The critical black hole mass is quite a bit larger than that found with the small clusters from Moore (1993) because M is larger. Note that $M_{\text{bh,crit}} \sim M/10$ for this cluster, which is on the verge of the limit in which many black holes will be inside the cluster at any given time.

Klesson & Burkert (1995) recently argued that Moore’s determination of $M_{\text{bh,crit}}$ for his set of nine diffuse clusters was flawed because deviations from the mean energy input for each encounter history would be quite large, implying a significant scatter in the values of $M_{\text{bh,crit}}$. They contend that clusters are destroyed as a consequence of a small number of encounters with large energy inputs. Instead, we find that for eight of Moore’s nine clusters, all except PAL 4, disruption is due to numerous encounters with $b \lesssim r_{\text{core}}$ which cause small changes individually. For these light clusters, the expected number of impacts, $N_{\text{core}}(M_{\text{bh,crit}})$, inside r_{core} is large for $M_{\text{bh}} = M_{\text{bh,crit}}$. We find that five of Moore’s clusters have $N_{\text{core}}(M_{\text{bh,crit}}) > 200$, two have $N_{\text{core}}(M_{\text{bh,crit}}) > 50$, one has $N_{\text{core}}(M_{\text{bh,crit}}) = 20$, and PAL 4 has $N_{\text{core}}(M_{\text{bh,crit}}) \lesssim 1$. So (aside from PAL 4) the






Figure 8. Fraction of clusters destroyed in $T = 10^{10}$ years plotted against black hole mass. These curves were generated using the Gaussian Monte Carlo method.

expected fluctuations in the $N_{\text{core}}(M_{\text{bh,crit}})$ are $\lesssim 20$ per cent for eight of Moore’s clusters, and $\lesssim 10$ per cent for seven of them. Only for PAL 4, which sits at a large galactocentric radius where few encounters occur, is the evolution near $M_{\text{bh,crit}}$ very stochastic. This case will be discussed in the next section.

7 THE LARGE M_{bh} LIMIT FOR THE SURVIVAL PROBABILITY

7.1 Theory

The previous section treated the small M_{bh} regime in which many collisions slowly add energy and remove mass from a cluster. This approximation will be valid for $M_{\text{bh}} \ll M_{\text{high}}$, the mass at which the cluster can be disrupted in a single collision. In the opposite limit for which $M_{\text{bh}} \gtrsim M_{\text{high}}$, any collision within a ‘destructive impact parameter’, b_d , will destroy the cluster. In addition, the cumulative effect of many collisions outside b_d can also destroy the cluster.

Formulae for the changes in E , M , and ν for a single collision in the tidal limit were derived in §5.3; see eq. (37), (38), and (44). The three formulae have the same scaling with η_c and b and only differ by a factor which depends on $\tilde{\psi}_0$. Hence, in the following derivations we only use the formula for energy input to derive results explicitly. To find the expressions for disruption by mass loss or change in ν one must only substitute $C_E \rightarrow C_M, C_\nu$.

The energy input can be rewritten as

$$\frac{\Delta E}{|E|} = C_E \eta_c^2 \frac{r_t^4}{b^4} \equiv f \left(\frac{b_d}{b} \right)^4 \quad (59)$$

where the impact parameter for $\Delta E/|E| = f$ is

$$b_d = r_t \eta_c^{1/2} \left(\frac{C_E}{f} \right)^{1/4}. \quad (60)$$

This energy input is sharply peaked about the mean for $b \simeq b_d$ because the impact parameter at which $\Delta E \simeq \sigma_{\Delta E}$ is larger than b_d by a factor of $N^{1/2}$.

A cluster has been destroyed if an impact occurs with $b < b_d$. The critical impact parameter $b_d \geq r_t$ for $M_{bh} \geq M_{high} = (M_{Vrel}/\sigma_0)(f/C_E)^{1/2}$. Using $\langle V_{rel} \rangle \simeq 1.47V_c$ [see eq. (17)], and $\sigma_0 \simeq 1 - 10 \text{ km s}^{-1}$ implies $M_{high} \sim 10 - 100M$.

The mean number of destructive encounters, N_d , which occur in a time T is given by the integrated rate of encounters inside $b_d(V_{rel})$:

$$N_d(T) = \pi n_{bh} T \int_0^\infty dV_{rel} f(V_{rel}) V_{rel} b_d^2(V_{rel}) = \left(\frac{C_E}{f} \right)^{1/2} \frac{\rho_{bh}}{M} \pi r_t^2 \sigma_0 T. \quad (61)$$

Note that $N_d(T) \propto f_{halo}$ but is independent of M_{bh} . Consequently, when $M_{bh} > M_{high}$, limits can be derived for f_{halo} but not M_{bh} (Wielen 1988).

For given f and T , the Poisson probability of no individually destructive events is given by

$$P_s(f_{halo}) = \exp[-N_d(f_{halo})]. \quad (62)$$

The expression for P_s becomes more complicated when we include the cumulative effect of all the nondestructive encounters outside b_d . As most collisions have $b \gg b_d$, they do not significantly change the cluster properties. These gentle collisions can be modelled using diffusion coefficients.

The mean rate of change of E due to impacts outside b_d is given by

$$\frac{\dot{E}}{|E|} = n_{bh} \int_0^\infty dV_{rel} f(V_{rel}) V_{rel} \int_{b_d(V_{rel})}^\infty 2\pi b db \left[f \frac{b_d^4(V_{rel})}{b^4} \right] = f \Gamma_d \quad (63)$$

where $\Gamma_d = N_d(t)/t$. Similarly, the time derivative of the variance is given by

$$\frac{\dot{\sigma}_E^2}{E^2} = n_{bh} \int_0^\infty dV_{rel} f(V_{rel}) V_{rel} \int_{b_d(V_{rel})}^\infty 2\pi b db \left[f^2 \frac{b_d^8(V_{rel})}{b^8} \right] = \frac{1}{3} f^2 \Gamma_d \quad (64)$$

where we have ignored the small variance per collision; this variance is due only to the variable number of encounters in a given time period. In addition, collisions inside b_d act as a ‘sink’ for clusters. The appropriate equation for $P(E, t)$, the probability density at time t and energy E , is

$$\frac{\partial P}{\partial t} = -\Gamma_d P - \dot{E} \frac{\partial P}{\partial E} + \frac{1}{2} \sigma_E^2 \frac{\partial^2 P}{\partial E^2} \quad (65)$$

where again the diffusion coefficients are treated as constants for simplicity. The solution with $E = E_0$ at $t = 0$ is

$$P(E, t) = \frac{\exp[-\Gamma_d t]}{(2\pi \dot{\sigma}_E^2 t)^{1/2}} \exp \left[-\frac{(E - E_0 - \dot{E} t)^2}{2 \dot{\sigma}_E^2 t} \right]. \quad (66)$$

This expression reduces to the $M_{bh} < M_{high}$ solution when $b_d \rightarrow 0$. The probability of survival is then the cumulative probability that the change in E has been less than $f|E|$, or

$$P_s = \frac{1}{2} e^{-\Gamma_d T} \left[1 + \text{erf} \left(\frac{f|E| - \dot{E} T}{\sqrt{2 \dot{\sigma}_E^2 T}} \right) \right] = \frac{1}{2} e^{-N_d} \left[1 + \text{erf} \left(\frac{1 - N_d}{\sqrt{2 N_d / 3}} \right) \right]. \quad (67)$$

For small N_d , $P_s \simeq \exp(-N_d)$, as in eq. (62), but for large N_d , $P_s \simeq (6\pi N_d)^{-1/2} \exp(-5N_d/2)$, which decreases much faster. Analogous results can be derived for changes in ν and M .

This decrease in the survival probability below the simple exponential decline due to the effects of distant encounters has been noted by previous investigators. In Wielen (1988), Fig. 5 presented a numerical determination of the survival probability in the large M_{bh} limit for clusters bombarded by both black holes and giant molecular clouds. Using $N_d(T) \simeq 0.5 \times T/T_{1/2}$, where $T_{1/2}$ is the time over which half the clusters have dissolved, our eq. (67) is a rather good fit to the figure.

The results of this section do not agree with Klesson & Burkert (1995) since the curves of P_s against M_{bh} in their Figs. 8, 9, and 10 show no systematic evidence of an asymptotic P_s independent of M_{bh} . This is surprising as they have included exponential adiabatic damping [Spitzer (1958), but see also Murali & Arras (1998) and Weinberg (1994) for a more recent

Table 4. Limits on f_{halo} in the Large M_{bh} Limit

Cluster Name	Criterion ^(a)	$M_{\text{high}}/M_{\odot}$	$f_{\text{halo}}(P_s = 0.9)$	$f_{\text{halo}}(P_s = 0.5)$	$f_{\text{halo}}(P_s = 0.1)$
AM 4	ν	3.8×10^5	0.028	0.14	0.33
ARP 2	ν	1.6×10^7	0.014	0.07	0.16
NGC 5053	ν	1.8×10^7	0.011	0.06	0.13
NGC 7492	ν	1.5×10^7	0.029	0.15	0.34
PAL 4	ν	1.4×10^7	0.174	0.90	>1.0
PAL 5	ν	8.6×10^6	0.005	0.02	0.05
PAL 13	ν	6.4×10^6	0.053	0.27	0.63
PAL 14	ν	8.7×10^6	0.035	0.18	0.41
PAL 15	ν	4.4×10^6	0.025	0.13	0.30
AM 4	20%M	4.1×10^5	0.031	0.20	0.67
ARP 2	20%M	7.6×10^6	0.006	0.04	0.14
NGC 5053	20%M	1.0×10^7	0.006	0.04	0.13
NGC 7492	20%M	4.9×10^6	0.009	0.06	0.20
PAL 4	20%M	8.9×10^6	0.109	0.72	>1.0
PAL 5	20%M	5.8×10^6	0.003	0.02	0.07
PAL 13	20%M	2.1×10^6	0.018	0.12	0.39
PAL 14	20%M	6.1×10^6	0.025	0.16	0.54
PAL 15	20%M	3.9×10^6	0.023	0.15	0.50

(a) The criterion ‘ ν ’ means that neither single destructive events nor many nondestructive events caused $\nu \rightarrow -0.6$. The criterion ‘20%M’ means that no single events of 20 per cent mass loss occurred.

version of these effects] which would decrease the energy input per encounter and hence *increase* P_s . The lack of encounters with small energy input in Klesson & Burkert’s (1995) simulations could result if their method of choosing the collision parameters oversampled small b and V_{rel} compared to our method.

Table 4 contains the allowed values of f_{halo} derived from the fixed cluster approximation of this section for the nine clusters found in Moore (1993). Two different destruction criteria were used: (1) ν out of bounds in $T = 10^{10}$ years due to the combined influence of single destructive collisions and multiple nondestructive collisions [eq. (67)], and (2) a single episode of 20 per cent mass loss in $T = 10^{10}$ years [eq. (62)]. Also given are the values of M_{high} , the black hole mass above which the cluster can be destroyed in a single collision, appropriate to each criterion. Three values were used for the probability of survival: $P_s = 0.1, 0.5$, and 0.9 .

For 50 per cent survival probability using the criterion $\nu \rightarrow -0.6$, the limiting values of f_{halo} range from 0.02 (PAL 5) to 0.90 (PAL 4). It is unlikely that $f_{\text{halo}} > 0.3$ since $P_s < 0.1$ for most of the clusters in that case. These limits depend sensitively on galactocentric radius and cluster size, as is evident in the scatter in the values of f_{halo} .

The results of this section will be tested in §8 using the results of the full Monte Carlo simulations. One wrinkle which appears in the results is that the different criteria for destruction can compete with each other so that the results of this section are not always good approximations to $P_s(N_d)$. Only in the cases where one method of destruction completely dominates over all the others do the results agree accurately.

8 MONTE CARLO SIMULATIONS OF INDIVIDUAL COLLISIONS

8.1 The Set-up

Previously, $M_{\text{bh,crit}}$ and f_{halo} were calculated for the two limiting cases of slow heating (small M_{bh}) and single-event destruction (large M_{bh}). The major simplification for both methods was the neglect of detailed, collision-by-collision evolution of the cluster. Both calculations relied on approximations. The Gaussian Monte-Carlo method relied on restricting collisions to very small energy and mass changes individually, and involved averaging over many collisions to streamline the computations. In this section we present a Monte-Carlo simulation of individual black hole encounters. We make no approximations for the energy input besides the impulse and straight line approximations; consistent with the former, we neglect displacement of cluster stars during the encounter. This treatment is simpler than an N-body simulation because the evolution of the cluster is mapped by the sequence of King models. As before, the King sequence limits the evolution we allow.

The calculations amount to simulating the ‘Green’s function’ of the cluster. A cluster in a given initial configuration is subjected to collisions with the halo black hole population. After each collision the King model for the cluster is altered, using its post-collision E and M , and assuming $r_t \propto M^{1/3}$. An evolutionary history of the cluster is mapped out over 10^{10} years,

and a final cluster state is found. As this process depends on many random variables, a range of final states is possible for each initial state, and many realizations are needed to find the Green's function.

8.2 The Probability Distributions for t , b , and V_{rel}

In §4, we derived the distribution of relative cluster-black hole speeds assuming an isothermal black hole halo. For the Monte Carlo simulations, we need to know the probability that the *next* collision suffered by a cluster occurs a time interval t to $t + dt$ after a given encounter, and involves an impact parameter in $(b, b + db)$ and a relative speed in $(V_{\text{rel}}, V_{\text{rel}} + dV_{\text{rel}})$. Since the number of collisions inside b is $\propto b^2$, we chose a maximum impact parameter b_{max} , and define $\sigma_{\text{max}} = \pi b_{\text{max}}^2$. Let us first consider collisions with a single relative velocity, V_{rel} . The expected number of collisions in a time t is

$$\overline{N} = \int_0^t n_{\text{bh}} \sigma_{\text{max}} V_{\text{rel}} d\tau, \quad (68)$$

so the Poisson probability that there are no collisions for the time interval $(0, t)$ and then one collision in $(t, t + dt)$ is

$$dP_1(t) = \exp \left[- \int_0^t n_{\text{bh}} \sigma_{\text{max}} V_{\text{rel}} d\tau \right] \cdot n_{\text{bh}} \sigma_{\text{max}} V_{\text{rel}} dt. \quad (69)$$

Multiplying by the probability $2\pi b db / \sigma_{\text{max}}$ that the collisions is in $(b, b + db)$ gives

$$dP_1(t, b) = \left\{ \exp \left[- \int_0^t n_{\text{bh}} \sigma_{\text{max}} V_{\text{rel}} d\tau \right] \times n_{\text{bh}} \sigma_{\text{max}} V_{\text{rel}} dt \right\} \times \left\{ \frac{2\pi b db}{\sigma_{\text{max}}} \right\}. \quad (70)$$

The first bracketed factor can be used to choose t , while the second bracketed factor can be used to choose b .

Next, suppose there is a set of black hole populations, each with a single relative velocity $V_{\text{rel},i}$; let f_i be the fraction of all black holes with $V_{\text{rel},i}$. Then the probability that the next collision has relative speed $V_{\text{rel},i}$ is

$$dP_1(t, i, b) = \left\{ \exp \left[- \int_0^t n_{\text{bh}} \sigma_{\text{max}} \langle V_{\text{rel}} \rangle d\tau \right] n_{\text{bh}} \sigma_{\text{max}} \langle V_{\text{rel}} \rangle dt \right\} \times \left\{ \frac{f_i V_{\text{rel},i}}{\langle V_{\text{rel}} \rangle} \right\} \times \left\{ \frac{2\pi b db}{\sigma_{\text{max}}} \right\} \quad (71)$$

where $\langle V_{\text{rel}} \rangle = \sum_i f_i V_{\text{rel},i}$. Take the continuum limit by substituting $f_i \rightarrow f(V_{\text{rel}}) dV_{\text{rel}}$ to get

$$dP_1(t, b, V_{\text{rel}}) = \left\{ \exp \left[- \int_0^t n_{\text{bh}} \sigma_{\text{max}} \langle V_{\text{rel}} \rangle d\tau \right] n_{\text{bh}} \sigma_{\text{max}} \langle V_{\text{rel}} \rangle dt \right\} \times \left\{ \frac{f(V_{\text{rel}}) V_{\text{rel}} dV_{\text{rel}}}{\langle V_{\text{rel}} \rangle} \right\} \times \left\{ \frac{2\pi b db}{\sigma_{\text{max}}} \right\} \quad (72)$$

The values of t , V_{rel} , and b may be chosen from the distributions in the first, second, and third bracketed factors, respectively.

Note that our derivation does not require $\overline{\Gamma} \equiv n_{\text{bh}} \sigma_{\text{max}} \langle V_{\text{rel}} \rangle$ to be time independent, so eq. (72) holds for any orbit, not just circular ones. Also, the velocity distribution found in the middle brackets is *not* $f(V_{\text{rel}})$, but is the distribution of speeds measured for the first collision after some chosen reference time. For example, the mean of this distribution is

$$\overline{V_{\text{rel}}} = V_c \left[\frac{3/2 + x}{\left(x + \frac{1}{2x}\right) \text{erf}(x) + \frac{1}{\sqrt{\pi}} \exp(-x^2)} \right] \quad (73)$$

where $x = V_{\text{cl}}/V_c$. As $x \rightarrow 0$, $\overline{V_{\text{rel}}}/V_c \rightarrow 3\sqrt{\pi}/4$, and as $x \rightarrow \infty$, $\overline{V_{\text{rel}}}/V_c \rightarrow 1$; for circular orbits eq. (17) and eq. (73) imply $\overline{V_{\text{rel}}}(x=1)/\langle V_{\text{rel}}(x=1) \rangle \simeq 1.16$, or, $\overline{V_{\text{rel}}}(x=1) \simeq 1.70 \times V_c$.

For globular clusters on circular orbits, the rate of encounters, $\Gamma = n_{\text{bh}} \sigma_{\text{max}} \langle V_{\text{rel}} \rangle$ inside σ_{max} is constant. The normalized cumulative distribution for the collision time is then $P_{\text{cdf},t}(t) = 1 - e^{-\Gamma t}$.

8.3 Evolving the Cluster

Distinct histories are computed for individual clusters at fixed R_g with identical initial values of M , r_t , r_{core} , and N , the number of cluster stars. The mass per star is held fixed at $m = (M/N)|_{t=0}$ even as M (and N) evolve; multiple species with different particle masses and spatial distributions were not considered. The maximum impact parameter b_{max} is also held fixed.

At any time $t \geq 0$, the cluster has known values of M , r_t , and E which allow determination of $\tilde{\psi}_0$, and hence the smooth King model phase space distribution. From this distribution, we choose random positions and velocities for the N stars in the cluster at the time of its *next* encounter with a black hole, $t_{\text{next}} > t$ (to keep fluctuations to a minimum, we actually choose velocities first in symmetric pairs $\pm \mathbf{v}$ and then choose positions from their distribution given the velocity). The values of t_{next} , b , and V_{rel} are chosen according to eq. (72), and velocity kicks $\Delta \mathbf{v}_i$ are computed for each cluster star using eq. 18 (by definition $\mathbf{b} = b \mathbf{e}_x$ and $\mathbf{V}_{\text{rel}} = V_{\text{rel}} \mathbf{e}_z$).

Given the $\Delta \mathbf{v}_i$, the center of mass velocity kick is $\overline{\Delta \mathbf{v}} = (1/N) \sum_{i=1}^N \Delta \mathbf{v}_i$; even stars which are ejected are included in the sum. Star i is ejected from the cluster if $(\mathbf{v}_i + \Delta \mathbf{v}_i - \overline{\Delta \mathbf{v}})^2 > 2\psi(r_i)$, where $\psi(r_i)$ is the pre-collision potential at r_i . If N' stars remain in the cluster then $M' = M(N'/N)$ is the new mass of the cluster. The new energy of the cluster is the total energy input to the remaining stars ($i = 1, 2, \dots, N'$) plus the contribution from the ejected stars ($j = 1, 2, \dots, N - N'$);

$$\Delta E = \sum_{i=1}^{N'} m \left(\mathbf{v}_i \cdot \delta \mathbf{v}_i + \frac{1}{2} |\delta \mathbf{v}_i|^2 \right) + \sum_{j=1}^{N-N'} m \left(\frac{GM}{r_t} + \psi_j - \frac{1}{2} v_j^2 \right). \quad (74)$$

The new parameters r'_t and $\tilde{\psi}'_0$ are then found by the steps outlined in §3. In this way, we have a new nondimensional model of the cluster after the collision.

These steps are repeated until either $t = 10^{10}$ years or the cluster ‘dies’. We keep track of six different conditions for destruction. For two of these, the simulation is stopped at once. The first, ‘ ν out of bounds’, triggers if ν goes out of the range $(-2.13, -0.6)$ (since all clusters simulated have $\tilde{\psi}_0 \lesssim 5.5$), signalling that no unique member of the King sequence can be found to represent the cluster. In our simulations, clusters always went out of bounds at $\nu \rightarrow -0.6$, so below we refer to this condition as ‘ $\nu \rightarrow -0.6$ ’. The second criterion for an immediate halt, called ‘ M_{20} ’, is realized if a *single* event of $|\Delta M| \geq 0.20M$ occurs which would greatly distort the cluster and invalidate our approximation of small $\Delta M/M$. There are four other criteria which are recorded *the first time they occur*, but do not stop the simulation. These are: ‘ M_{10} ’, a single occurrence of 10 per cent mass loss; ‘ $\frac{1}{2}M$ ’, decrease of the mass of the cluster to half its original value; ‘ $\frac{1}{2}E$ ’, change of the energy of the cluster up or down by half of its original value; and ‘ $\frac{1}{2}E/M$ ’, change of the quantity E/M up or down by half of its original value. Survival of the cluster will be called criterion ‘ s ’. Note that the criteria M_{10} , $\frac{1}{2}E$, $\frac{1}{2}M$, and $\frac{1}{2}E/M$ can happen repeatedly before $\nu \rightarrow -0.6$, M_{20} , or s , but are only recorded the first time they occur. The two conditions $\frac{1}{2}E$ and $\nu \rightarrow -0.6$ correspond most closely to the criteria used by previous investigators, but since our simulations include mass loss the correspondence is not exact. It is possible for a cluster to be destroyed in less than 10^{10} years by M_{20} or $\nu \rightarrow -0.6$ even though $\frac{1}{2}M$, $\frac{1}{2}E$, M_{10} , or $\frac{1}{2}E/M$ might not have had a chance to occur yet. Moreover, M_{20} may occur before $\nu \rightarrow -0.6$. When analyzing the results, the competition among the various criteria must be kept in mind.

This simulation ignores a number of possibly important effects. The most restrictive approximation is the use of the King sequence to model the evolution of the cluster. Relatively small changes in mass and energy may lead to $\tilde{\psi}_0 \rightarrow 0$, so a normal cluster could have a lifetime larger than is found here. For example, a $M \sim 10^6 M_\odot$ cluster could lose 30 per cent of its binding energy and mass, but no longer be fit well by a King model; yet you would still have a $M \sim 7 \times 10^5 M_\odot$ cluster. Our treatment assumes that clusters become unstable to rapid dissolution when $\tilde{\psi}_0 \rightarrow 0$.

To narrow the focus to the effects of halo black holes, we have neglected the disk and bulge components of the galaxy; destructive effects such as disk shocking and collisions with molecular clouds have also been suppressed entirely. Internal evolution of the cluster and non-spherical galactic tidal fields are not treated. Clusters are kept at single R_g , so time dependence of the density of halo black holes and relative velocities along cluster orbits is neglected. Lastly, we will make fractional errors of order $\Delta M/M$ in our method.

Nevertheless, our model advances previous attempts to constrain properties of a hypothetical population of halo black holes via their effects on clusters. We include mass loss which, as was shown in §5.3, contributes significantly to cluster heating, sometimes slightly more than ‘Spitzer’ heating. Moreover, since the mean mass loss is comparable to the mean energy input, the evolution of $\nu \propto E/M^{5/3}$ is driven by both ΔE and ΔM . Here no simplifications of the energy input are employed except the impulse approximation and the straight line orbit approximation. Hence, the Monte Carlo method for finding $\Delta E(b)$ includes the important ‘shocking’ effect of the $\mathbf{v} \cdot \delta \mathbf{v}$. Lastly, the correct expression for the rate of collisions in eq. (16) is used. We also note that many criteria for the cluster to be disrupted have been used in the past; the most popular is $\Delta E(t)/|E| = 1$. This choice is usually implemented with no regard to mass loss and evolution. Here, we test a variety of criteria for cluster destruction in order to find the most restrictive.

One last technical detail which must be discussed is the value of $b_{\max}$, the maximum impact parameter for our scattering experiment. Since the number of collisions that must be simulated $\propto b_{\max}^2$, it is essential to choose as small a $b_{\max}$ as possible without losing accuracy. First examine the small M_{bh} case in which the cluster cannot be destroyed in a single pass. In this slow heating limit, $\Delta E(b) \propto b^{-4}$ outside the core and $\int_0^{b_{\max}} 2\pi b db \Delta E(b)$ converges as $b_{\max}^{-2}$. Nearly all of the heating results from penetrating encounters and $b_{\max} \simeq r_t$ will be a good approximation. However, in the large M_{bh} case there is a critical impact parameter $b_d^2 \propto M_{\text{bh}}$ inside of which the cluster is destroyed, and we must choose $b_{\max} > b_d \propto M_{\text{bh}}^{1/2}$. An estimate for M_{high} , the value of M_{bh} above which $b_{\max}$ must increase $\propto M_{\text{bh}}$, is discussed in §7.

8.4 Description of the Simulations

The time necessary to realize a cluster with $N = 1000$ stars and compute the energy input was ~ 1 sec on a Sun workstation. The computation time needed to map out the history for one cluster is $\overline{N_{ev}} \times 1 \text{ sec} \sim (80 \text{ hrs})(M/M_{\text{bh}})(10 \text{ pc}/r_t)$. This severely

limited practical choices for N and m . In order for the simulations to be realistic, we chose to simulate the full range of M_{bh} for AM 4 only, using $m = 0.7M_{\odot}$ and $N = 1000$ initially.

To study much more massive clusters, the time limitation would force the number of stars used to be a small fraction of the physical value. As a consequence, the variance in energy input and mass loss would be unrealistically large in the simulations, and hence the Green's function would be spread over too broad a range of final states. Too small a number of stars will decrease the number of stars in the portion of phase space from which particles are ejected resulting in an incorrect evaluation of mass loss, especially if the total mass lost in the simulation is small. Uneven sampling of phase space, arranged with finer sampling near the escape surface, could alleviate this problem.

For a given M_{bh} , b_{max} , and f_{halo} , a number N_t trials were performed. The number of clusters destroyed, N_{dest} , by each of the six criteria was recorded and the fraction $f_{\text{dest}} = N_{\text{dest}}/N_t$ of clusters destroyed computed. In addition to f_{dest} for each criterion, the sum of $f_{\text{dest}}(\nu \rightarrow -0.6) + f_{\text{dest}}(M_{20})$ is computed because the sum of the fraction destroyed by these two conditions is not subject to competition effects which appear for the six criterion separately.

At least two runs with different b_{max} were done for each cluster at a certain M_{bh} . The runs with the larger b_{max} are presented here. The variation in f_{dest} for the two runs was in all cases within the error bars shown in the figures. The value of b_{max} used for the results presented here was $b_{\text{max}} = 2r_t$ for $M_{\text{bh}} \lesssim 100M$ and $b_{\text{max}} = 2r_t(M_{\text{bh}}/100M_{\odot})^{1/2}$ for $M_{\text{bh}} > 100M$. These values of b_{max} are much larger than is needed, as $M_{\text{high}} \gtrsim 4 \times 10^5 M_{\odot} = 570M$ (see Table 3), and the destructive radius is not outside r_t until $M_{\text{bh}} > M_{\text{high}}$.

The number, N_t , of trials ranged between $N_t = 40$ for small M_{bh} and $N_t = 1000$ for large M_{bh} . The number of trials was restricted by the run time, which was greatly increased for small M_{bh} because of the large number of collisions $\propto f_{\text{halo}}/M_{\text{bh}}$. Bayesian methods were used to compute P_s and its uncertainty. The error bars in the figures span the range of P_s around the peak of its posterior containing 68 per cent probability.

8.5 Results and Discussion

The first set of runs was for $f_{\text{halo}} = 1$. The fractions of trial clusters destroyed by our various criteria are plotted in Fig. 9.

The dominant destruction mechanism by far is $\nu \rightarrow -0.6$. For this mechanism, the black hole mass at which 50 per cent of the clusters were destroyed is

$$M_{\text{bh,crit}}(50\%) \simeq 600M_{\odot} = 0.86M \quad (75)$$


The other destruction mechanisms only come into play once $M_{\text{bh}} \gtrsim 5 \times 10^4 M_{\odot}$. At this point, $\nu \rightarrow -0.6$ still destroys 100 per cent of the clusters (as predicted in Table 4 for $f_{\text{halo}} = 1.0$); even the approximately 40 per cent that die via M_{20} do so in the same encounter that pushes $\nu \rightarrow -0.6$. If $\nu \rightarrow -0.6$ were not stopping the simulations the other destruction criteria would be more important, and f_{dest} would not necessarily asymptote to small values as in Fig. 9.

We can compare the results in Fig. 9 for the criterion $\nu \rightarrow -0.6$ with the Gaussian Monte-Carlo simulations in Fig. 8. There the critical black hole mass is $M_{\text{bh,crit}}(50\%) \simeq 650M_{\odot}$, which is different from the full Monte-Carlo simulation by $\lesssim 10$ per cent. Considering that the error bars in f_{dest} are of order 10 – 15 per cent, the two numbers agree. In addition, the mass range for which $0.9 \geq f_{\text{dest}} \geq 0.1$ is $\delta M_{\text{bh,crit}} \sim 250M_{\odot}$ for both the Gaussian Monte-Carlo and full Monte-Carlo cases. The agreement of these two methods shows that the assumptions of (1) a Gaussian distribution of energy input and mass loss, and (2) smooth Fokker-Planck evolution, are accurate in the small M_{bh} regime. In addition, the ansatz for the variance of the number of stars ejected in §5.1 gave similar results as the full Monte-Carlo treatment, as evidenced by the comparable widths $\delta M_{\text{bh,crit}}$.

The second set of runs explores the large M_{bh} limit of §7. Two runs for AM 4 at $M_{\text{bh}} = 1000M = 7 \times 10^5 M_{\odot}$ and $M_{\text{bh}} = 5000M = 3.5 \times 10^6 M_{\odot}$ were performed as discussed in the previous section. The results are shown in Fig. 10 and 11. In addition, we also tried NGC 5053 since it has a fairly large mass, $M = 37700M_{\odot}$, and yet it restricts f_{halo} to be small. Because the run time would have been prohibitively long using the correct number of stars in this cluster, we chose $N = 1000$ for NGC 5053. As discussed previously, this should not change the values of P_s much since we will be in the limit in which the collisions are quite strong, and the variances are expected to be small compared to the means. A single black hole mass of $M_{\text{bh}} = 1000M = 3.77 \times 10^7 M_{\odot}$ was used to get the results shown in Fig. 12.

The values of f_{halo} used in Fig. 10 and 11 are (from left to right), $f_{\text{halo}} = 0.05, 0.1, 0.2, 0.3, 0.4, 0.5, 0.6$. The values of N_d computed with eq.(61) for each cluster will depend on the criterion for destruction through the values of f and C chosen. This is why the two curves for the two criteria $\nu \rightarrow -0.6$ and $\Delta M/M = 20$ per cent do not have the same values of N_d for each data point of a given f_{halo} . We have converted the values of f_{halo} to the values of N_d shown in the graphs using eq. (60) and (61). The values of f_{halo} used for NGC 5053 in Fig. 12 are (from left to right), $f_{\text{halo}} = 0.01, 0.055, 0.128, 0.17, 0.273$.

As the results for AM 4 and NGC 5053 were a bit different, let us discuss AM 4 first. Notice that the full Monte-Carlo P_s curves in Figs. 10 and 11 agree quite well with each other. This was predicted in §7 from the fact that N_d , the number of destructive encounters, is independent of M_{bh} . Next notice that the curves for the $\nu \rightarrow -0.6$ destruction criterion agree quite well with eq. (67) which was used to draw the solid lines in the figures. The predicted P_s for $\Delta M/M = 20$ per cent did






Figure 9. Fraction of initial AM 4 clusters destroyed according to the various criteria for destruction as a function of M_{bh} . These curves were made using the full Monte Carlo method.




Figure 10. Comparison of full Monte Carlo simulations against theory for AM 4 with $M_{\text{bh}} = 5000M$. The circles are the points of P_s vs. N_d for the destruction criterion $\nu \rightarrow -0.6$. The triangles are for P_s vs. N_d with the destruction criterion of 20 per cent mass loss in a single encounter.

not, however, agree very well with eq. (62). This can be attributed to the fact that $\nu \rightarrow -0.6$ generally occurs well before any collision with $\Delta M/M = 20$ per cent. In fact, the good agreement of the numerical results with eq. (67) for $\nu \rightarrow -0.6$ is partly a consequence of the relative rarity of collisions with $|\Delta M|/M = 0.2$ for AM 4.

For NGC 5053, the fraction of clusters destroyed by $\Delta M/M = 20$ per cent is much larger, and hence neither curve agrees well with either eq. (67) or (62). For $N_d \lesssim 1.5$, since few clusters were destroyed by $\Delta M/M = 20$ per cent the agreement for $\nu \rightarrow -0.6$ was good, but competition between the two destruction criteria is apparent for $N_d \gtrsim 1.5$. However, note that the survival probability against $\nu \rightarrow -0.6$ is smaller than for $|\Delta M|/M = 0.2$. Since the latter occurs *only* in a single catastrophic encounter, whereas the former also involves the cumulative effect of numerous gentle collisions, it is incorrect to ascribe cluster destruction at large M_{bh} entirely to the effect of the single most destructive collision.




Figure 11. Comparison of full Monte Carlo simulations against theory for AM 4 with $M_{bh} = 1000M$. The circles are the points of P_s vs. N_d for the destruction criterion $\nu \rightarrow -0.6$. The triangles are for P_s vs. N_d with the destruction criterion of 20 per cent mass loss in a single encounter.




Figure 12. Comparison of full Monte Carlo simulations against theory for NGC 5053 with $M_{bh} = 1000M$. The circles are the points of P_s vs. N_d for the destruction criterion $\nu \rightarrow -0.6$. The triangles are for P_s vs. N_d with the destruction criterion of 20 per cent mass loss in a single encounter.

9 CONCLUSIONS AND PROSPECTS FOR FUTURE WORK

Our results confirm the basic picture of cluster evolution proposed by Wielen (1988) in which there are two distinct regimes based on the size of the black hole mass; for M_{bh} too small to destroy the cluster in a single collision, the evolution can be modelled by a smooth average over many collisions while for M_{bh} large enough to destroy the cluster in a single collision, the survival probability over 10^{10} years depends both on the heating from non-destructive encounters and the stochastic effect of the individually destructive collisions.

Several technical improvements have been made over previous investigations for the $M_{bh} \ll M_{high}$ limit. The cluster's structure was allowed to change by evolving it along a King sequence; comparison of the values of $M_{bh,crit}$ in Fig. 8 and the last column of Table 3 show that evolution accelerates the dissolution process for Moore's weakly bound clusters, leading to a stricter limit. The inclusion of mass loss gave energy changes comparable to the usual $\delta v^2/2$ heating. Finite N effects have been included, and shown to be relatively unimportant in determining $M_{bh,crit}$ for the $M_{bh,crit} \ll M_{high}$ case since 'the range

of final states' is quite small when the cluster is evolved over times long enough to disrupt it. The Fokker-Planck model (§6.2) for the evolution of an ensemble of initial clusters agreed closely with the full Monte-Carlo simulations (§8) showing that the 'slow heating' approximation is indeed an accurate representation of the evolution. Indeed, the simple model of §6.2 with constant diffusion coefficients given by the convenient formulas in Table 2 gives quick, analytical results accurate to within a factor of a few (Table 3).

The final fate of the clusters studied in this paper was always dissolution. This occurred for three reasons. First, we ignored internal relaxation which tends to drive a cluster to core collapse. The role of internal relaxation is currently being studied (Murali et al. 1998). Second, when the cluster is heated slowly by many penetrating encounters, the final fate is dissolution independent of cluster concentration. Third, in the large M_{bh} limit in which all non-destructive encounters are tidal, clusters with $\tilde{\psi}_0 \lesssim 5.5$ dissolve and those with $\tilde{\psi}_0 \gtrsim 5.5$ core collapse. All the clusters examined here had $\tilde{\psi}_0 \lesssim 5.5$.

Our results for the slow-heating, $M_{bh} \ll M_{high}$ limit are given in Table 3, Fig. 8, and Fig. 9. The strictest limit on M_{bh} comes from the full Monte-Carlo calculation for AM 4 with $M_{bh,crit} \simeq 600M_\odot$. For the ν out of bounds criterion, several of Moore's clusters are disrupted at $M_{bh} \sim 1000M_\odot$ and seven of nine are disrupted for $M_{bh} \sim 6000M_\odot$. For the $\Delta E/|E| = 0.5$ criterion, six clusters die at $M_{bh} \sim 5000$ and two die for $M_{bh} \sim 1500$. To summarize, in this regime it is extremely unlikely that all the clusters could survive unscathed for 10^{10} years if M_{bh} is greater than a few thousand solar masses.

For the $M_{bh} \gg M_{high}$ limit, the probability of survival P_s does not tend to zero as $M_{bh} \rightarrow \infty$, but instead asymptotes at a nonzero value (which does not have to be small). A simple Fokker-Planck model to determine P_s has been developed (§7) in which both the close, destructive collisions and the distant, non-destructive collisions are included. The distribution of energies for an initial ensemble of clusters is shown to depend only on the expected number of destructive encounters for the cluster as a function of time, and the resultant P_s gives good agreement with the full Monte-Carlo simulations (when the competition of the various survival criteria is small). Inclusion of the distant, non-destructive encounters leads to smaller values of P_s which in turn gives much tighter limits on the allowed fraction of the halo in black holes.

The results for the $M_{bh} \gg M_{high}$ limit are presented in Table 4 and Fig. 9, 10, 11, and 12. For a particular cluster and a given value of P_s , one can place limits on f_{halo} . For $P_s = 0.5$, Table 4 gives values of f_{halo} for Moore's clusters ranging from 0.02 to 0.9. In this "tidal" regime, it is unlikely that $f_{halo} > 0.3$ since then $P_s < 0.1$ for most of Moore's clusters.

The existence for 10^{10} years of the set of nine tenuous globular clusters studied by Moore (1993) places severe restrictions the allowed mass and fraction of the halo mass in massive black holes. Left unanswered in our paper is the question of whether Moore's clusters were *initially* similar in size to what we see today, or if they have been 'whittled down' to their present small stature. Indeed, even in isolation, the least massive of these clusters, AM 4, PAL 13, and perhaps NGC 7492 might be expected to have evaporated in $\sim 10^9$ years, suggesting that they are relatively young (Murali 1998; Murali et al. 1998; Spitzer 1987). To answer this question fully, one must evolve a representative population of *initial* clusters and attempt to reproduce the observed population today. The work for this project has already begun (Murali et al. 1998). Significantly tighter limits may result from these new investigations by the introduction of other sources of cluster evolution, such as internal relaxation, evaporation, and disk shocking.

ACKNOWLEDGMENTS

We would like to thank David Chernoff and Chigurupati Murali for many informative discussions, Mark Scheel for invaluable computer assistance, Jamie Lombardi for helpful comments, and Peter Brooks for the title. This research was supported in part by NSF grants no. AST 93-15375 and AST 95-30397.

REFERENCES

- Bahcall J.N., Hut P., Tremaine S., 1985, ApJ, 290, 15
- Binney J., Tremaine S., 1987, Galactic Dynamics. Princeton University Press, Princeton
- Carlberg R.G., Dawson P.C., Hsu T., VandenBerg D.A., 1985, ApJ, 294, 674
- Carr B., 1994, ARA&A, 32, 531
- Carr B., Sakkellariadou M., 1998, preprint
- Chernoff D.F., Kochanek C. S., Shapiro S. L., 1986, ApJ, 309, 183
- Gomez A.E., Delhaye J., Grenier S., Jaschek C., Arenou F., Jaschek M., 1990, A&A, 236, 95
- King I.R., 1966, AJ, 71, 64
- Klessen R., & Burkert A., 1995, MNRAS, 280, 735
- Lacey C.G., 1991, in Sundelius B., ed., Dynamics of Galaxies, Goteborg, pg.257
- Loeb A., 1993, ApJ, 403, 542
- Loeb A., Rasio F., 1994, ApJ, 432, 52
- Moore B. 1993, ApJ, 413, L93
- Murali C., 1998, personal communication.
- Murali C., Arras P., in preparation

- Murali C., Arras P., Wasserman I., in preparation
 Ostriker J.P., Lacey C.G., 1985, ApJ, 299, 633
 Ostriker J.P., Spitzer L., Chevalier R.A., 1972, ApJ, 193, L1
 Retterer J.M., King I.R., 1982, ApJ, 254, 214
 Spitzer L., 1958, ApJ, 127, 17
 Spitzer L., 1987, Dynamical Evolution of Globular Clusters. Princeton University Press, Princeton
 Spitzer L., and Chevalier R.A., 1973, ApJ, 183, 565
 Strömgren, B., 1987, in Gilmore G., Carswell B., eds, Proc. of NATO ASI, The Galaxy. Reidel, Cambridge, p.229
 Umemura M., Loeb A., Turner E.L., 1993, ApJ, 419, 459
 Wasserman I., Salpeter E., 1994, ApJ, 433, 670
 Wasserman I., Weinberg M.D., 1991, ApJ, 382, 149
 Weinberg M.D., 1994, AJ, 108(4), 1403
 Weinberg M.D., Shapiro S.L., Wasserman I., 1987, ApJ, 312, 367
 Weilen R., 1977, A&AS, 60, 263
 Weilen R., 1985, in Goodman J., Hut P., eds, Proc. IAU Symp. 113, Dynamics of Star Clusters. Reidel, Dordrecht
 Wielen R., 1988, in Grindlay J.E., Philip A.G.D., eds, Proc. IAU Symp. 126, The Harlow-Shapley Symposium on Globular Cluster Systems in Galaxies. Kluwer, Dordrecht, p.393
 Wiyanto P., Kato S., Inagaki S., 1985, PASJ, 37(4), 715

APPENDIX A: THE CENTER OF MASS VELOCITY KICK

For a continuous cluster with spherical mass density $\rho(r)$, the velocity kick to the cluster center of mass takes on a very simple form. For $\mathbf{b} = b\mathbf{e}_x$, $\mathbf{V}_{\text{rel}} = V_{\text{rel}}\mathbf{e}_z$, and projected cluster position $\mathbf{R} = R\cos(\phi)\mathbf{e}_x + R\sin(\phi)\mathbf{e}_y$, eq. (18) gives the mean velocity kick

$$\begin{aligned}
 \langle \Delta \mathbf{v} \rangle &= \frac{2GM_{\text{bh}}}{V_{\text{rel}}} \frac{1}{M} \int d^3x \rho(r) \frac{(b - R\cos(\phi))\mathbf{e}_x - R\sin(\phi)\mathbf{e}_y}{b^2 + R^2 - 2bR\cos(\phi)} \\
 &= \frac{2GM_{\text{bh}}}{V_{\text{rel}}} \frac{1}{M} \int_0^{r_t} dR R \Sigma(R) \int_0^{2\pi} d\phi \frac{(b - R\cos(\phi))\mathbf{e}_x - R\sin(\phi)\mathbf{e}_y}{b^2 + R^2 - 2bR\cos(\phi)} \\
 &= \frac{2GM_{\text{bh}}}{V_{\text{rel}}} \frac{1}{M} \int_0^{r_t} dR 2\pi R \Sigma(R) \frac{\theta(b - R)}{b} \mathbf{e}_x \\
 &= \frac{2GM_{\text{bh}}}{bV_{\text{rel}}} \frac{M_{\text{cyl}}(b)}{M} \mathbf{e}_x
 \end{aligned} \tag{A1}$$

where $\Sigma(R) = 2 \int_0^{(r_t^2 - R^2)^{1/2}} dz \rho(\sqrt{R^2 + z^2})$ and $M_{\text{cyl}}(b) = \int_0^b dR 2\pi R \Sigma(R)$ is the mass enclosed within a cylindrical distance b from the center of the cluster. The b dependent function $M_{\text{cyl}}(b)/bM$ is zero at the cluster center, reaches a maximum inside the cluster and is equal to $1/b$ outside the cluster since $M_{\text{cyl}}(b \geq r_t) = M$, recovering the usual velocity kick for a structureless particle.